

Homogenization and Asymptotics for Small Transaction Costs: The Multidimensional Case

DYLAN POSSAMAÏ¹, H. METE SONER^{2,3}, AND
NIZAR TOUZI⁴

¹CEREMADE, Université Paris Dauphine, Paris, France

²ETH (Swiss Federal Institute of Technology), Zurich, Switzerland

³Swiss Finance Institute, Zurich, Switzerland

⁴CMAP, Ecole Polytechnique Paris, Paris, France

In the context of the multi-dimensional infinite horizon optimal consumption investment problem with small proportional transaction costs, we prove an asymptotic expansion. Similar to the one-dimensional derivation in our accompanying paper, the first order term is expressed in terms of a singular ergodic control problem. Our arguments are based on the theory of viscosity solutions and the techniques of homogenization which leads to a system of corrector equations. In contrast with the one-dimensional case, no explicit solution of the first corrector equation is available and we also prove the existence of a corrector and its properties. Finally, we provide some numerical results which illustrate the structure of the first order optimal controls.

Keywords Asymptotic expansions; Homogenization; Transaction costs; Viscosity solutions.

2000 AMS Subject Classification 91B28; 35K55; 60H30.

1. Introduction

We continue our asymptotic analysis of problems with small transaction costs using the approach developed in [40] for problems with only one stock. In this paper, we consider the case of multi stocks with proportional transaction costs. The problem of investment and consumption in such a market was first studied by Magill and Constantinides [27] and later by Constantinides [9]. There is a large literature including the classical papers of Davis and Norman [11], Shreve and Soner [37] and Dumas and Luciano [13]. We refer to our earlier paper [40] and to the recent book of Kabanov and Safarian [24] for other references and for more information.

Received February 2, 2013; Accepted March 30, 2015

Address correspondence to H. Mete Soner, ETH (Swiss Federal Institute of Technology), Department of Mathematics, 101 Ramistrasse, Zurich, Switzerland; E-mail: hmsoner@ethz.ch

Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/lpde

This problem is an important example of a singular stochastic control problem. It is well known that the related partial differential equation contains a gradient constraint. As such it is an interesting free boundary problem. The main focus of this paper is on the analysis of the small transaction costs asymptotics. It is clear that in the limit of zero transaction costs, we recover the classical problem of Merton [29] and the main interest is on the derivation of the corrections of this obvious limit.

The asymptotic problem is a challenging problem which attracted considerable attention in the existing literature. The first rigorous proof in this direction was obtained in the appendix of [37]. Later several rigorous results [3, 5, 19, 23, 33] and formal asymptotic results [1, 20, 42] have been obtained. The rigorous results have been restricted to one space dimensions with the exception of the recent manuscript by Bichuch and Shreve [6]. As well known, utility indifference price in this market is an important approach as perfect hedging is very costly as shown in [39]. Davis et al. [12] were the first to study this approach with an exponential utility function and the formal asymptotics was later developed in [42].

In this paper, we use the techniques developed in [40]. As in that paper, the main technique is the viscosity approach of Evans to homogenization [14, 15]. This powerful method combined with the relaxed limits of Barles and Perthame [2] provides the necessary tools. As well known, this approach has the advantage of using only a simple L^∞ bound. In addition to [2, 14, 15], the rigorous proof utilizes several other techniques from the theory of viscosity solutions developed in the papers [2, 16, 17, 25, 32, 34, 38] for asymptotic analysis.

For the classical problem of homogenization, we refer to the reader to the classical papers of Papanicolaou and Varadhan [31] and of Souganidis [41]. However, we emphasize that the problem studied here does not even look like a typical homogenization problem of the existing literature. Indeed, the dynamic programming equation, which is the starting point of our asymptotic analysis, does not include oscillatory variables of the form x/ϵ or in general a fast varying ergodic quantity. The fast variable appears after a change of variables that includes the difference with the optimal strategy at the effective level problem. Hence, the fast variable actually depends upon the behavior of the limit problem, which is a novel viewpoint for homogenization where usually the fast variables are built into the equation from the beginning and one does not have to construct them from limits. This ergodic variable, in general, does not have to be periodic. In recent studies, deep techniques combining ergodic theorems and difficult parabolic estimates were used to study these more general cases. We refer the reader to Lions & Souganidis [26] for the almost-periodic case and Caffarelli and Souganidis [7] for the case in random media and to the references therein.

As in our accompanying paper [40], the formal asymptotic analysis leads to a system of corrector equations related to an ergodic optimal control problem similar to the monotone follower [4]. However, in contrast with the one-dimensional case studied in [40], no explicit solution is available for this singular ergodic control problem. This is the main difficulty that we face in the present multi-dimensional setting. We use the recent analysis of Hynd [21, 22] to analyze this multidimensional problem and obtain a $C^{1,1}$ unique solution of the corresponding “eigenvalue” problem satisfying a precise growth condition. This characterization is sufficient to carry out the asymptotic analysis. On the other hand, the regularity of the free boundary is a difficult problem and we do not study it in this paper. We refer the reader to [35, 36] for such analysis in a similar problem.

The paper is organized as follows. Section 2 provides a quick review of the infinite horizon optimal consumption-investment problem under transaction costs and recalls the formal asymptotics, as derived in [40]. Section 3 collects the main results of the paper. The rigorous proof of the first order expansion is reported in Section 4. In particular, the proof requires some wellposedness results of the first corrector equation which are isolated in Section 5. In the Appendix, we provide the proofs of several lemmas used in the proofs and report some numerical experiments which illustrate the nature of the first order optimal transfers.

Notations. Throughout the paper, we denote by \cdot the Euclidean scalar product in \mathbb{R}^d , and by (e_1, \dots, e_d) the canonical basis of \mathbb{R}^d . We shall be usually working on the space $\mathbb{R} \times \mathbb{R}^d$ with first component enumerated by 0. The corresponding canonical basis is then denoted by (e_0, \dots, e_d) . We denote by $\mathcal{M}_d(\mathbb{R})$ the space of $d \times d$ matrices with real entries, and by T the transposition of matrices. We denote by $B_r(x)$ the open ball of radius $r > 0$ centered at x , and $\bar{B}_r(x)$ the corresponding closure.

2. The General Setting

In this section, we briefly review the infinite horizon optimal consumption-investment problem under transaction costs and recall the formal asymptotics, derived in [40]. These calculations are the starting point of our analysis.

The next two subsections introduce the notations and known results that will be frequently used in our subsequent analysis. To be consistent with the earlier results, we report these subsections from [40].

2.1. Optimal Consumption and Investment under Proportional Transaction Costs

The financial market consists of a non-risky asset S^0 and d risky assets with price process $\{S_t = (S_t^1, \dots, S_t^d), t \geq 0\}$ given by the stochastic differential equations (SDEs),

$$\frac{dS_t^0}{S_t^0} = r(S_t)dt, \quad \frac{dS_t^i}{S_t^i} = \mu^i(S_t)dt + \sum_{j=1}^d \sigma^{i,j}(S_t)dW_t^j, \quad 1 \leq i \leq d,$$

where $r: \mathbb{R}^d \rightarrow \mathbb{R}_+$ is the instantaneous interest rate and $\mu: \mathbb{R}^d \rightarrow \mathbb{R}^d$, $\sigma: \mathbb{R}^d \rightarrow \mathcal{M}_d(\mathbb{R})$ are the coefficients of instantaneous mean return and volatility, satisfying the standard assumptions:

$$r, \mu, \sigma \text{ are bounded and Lipschitz, and } (\sigma\sigma^T)^{-1} \text{ is bounded.}$$

In particular, this guarantees the existence and the uniqueness of a strong solution to the above stochastic differential equations (SDEs).

The portfolio of an investor is represented by the dollar value X invested in the non-risky asset and the vector process $Y = (Y^1, \dots, Y^d)$ of the value of the positions in each risky asset. These state variables are controlled by the choices of the total amount of transfers $L_t^{i,j}$, $0 \leq i, j \leq d$, from the i -th to the j -th asset cumulated up to time t . Naturally, the control processes $\{L_t^{i,j}, t \geq 0\}$ are defined as càd-làg, nondecreasing, adapted processes with $L_{0-} = 0$ and $L^{i,i} \equiv 0$.

In addition to the trading activity, the investor consumes at a rate determined by a nonnegative progressively measurable process $\{c_t, t \geq 0\}$. Here c_t represents the rate of consumption in terms of the non-risky asset S^0 . Such a pair $v := (c, L)$ is called a *consumption-investment strategy*. For any initial position $(X_{0-}, Y_{0-}) = (x, y) \in \mathbb{R} \times \mathbb{R}^d$, the portfolio positions of the investor are given by the following state equation

$$dX_t = (r(S_t)X_t - c_t)dt + \mathbf{R}^0(dL_t), \quad \text{and} \quad dY_t^i = Y_t^i \frac{dS_t^i}{S_t^i} + \mathbf{R}^i(dL_t), \quad i = 1, \dots, d,$$

where

$$\mathbf{R}^i(\ell) := \sum_{j=0}^d (\ell^{j,i} - (1 + \epsilon^3 \lambda^{i,j}) \ell^{i,j}), \quad i = 0, \dots, d, \quad \text{for all } \ell \in \mathcal{M}_{d+1}(\mathbb{R}_+),$$

is the change of the investor's position in the i -th asset induced by a transfer policy ℓ , given a structure of proportional transaction costs $\epsilon^3 \lambda^{i,j}$ for any transfer from asset i to asset j . Here, $\epsilon > 0$ is a small parameter and the scaling ϵ^3 is chosen to state the expansion results simpler. Set

$$\mathcal{F} := \{(i, j) \in \{0, 1, \dots, d\}^2 : \lambda^{i,j} < \infty\}.$$

In the sequel, we always assume that $\lambda^{i,j} \geq 0$, $\lambda^{i,i} = 0$, for all i, j and $(0, i), (i, 0) \in \mathcal{F}$ for every $i = 0, 1, \dots, d$.

Let $(X, Y)^{v,s,x,y}$ denote the controlled state process. A consumption-investment strategy v is said to be *admissible* for the initial position (s, x, y) , if the induced state process satisfies the solvency condition $(X, Y)_t^{v,s,x,y} \in K_\epsilon$, for all $t \geq 0$, \mathbb{P} -a.s., where the solvency region is defined by:

$$K_\epsilon := \{(x, y) \in \mathbb{R} \times \mathbb{R}^d : (x, y) + \mathbf{R}(\ell) \in \mathbb{R}_+^{1+d} \text{ for some } \ell \in \mathcal{M}_{d+1}(\mathbb{R}_+)\}.$$

The set of admissible strategies is denoted by $\Theta^\epsilon(s, x, y)$. For given initial positions $S_0 = s \in \mathbb{R}_+^d$, $X_{0-} = x \in \mathbb{R}$, $Y_{0-} = y \in \mathbb{R}^d$, the consumption-investment problem is the following maximization problem,

$$v^\epsilon(s, x, y) := \sup_{(c,L) \in \Theta^\epsilon(s,x,y)} \mathbb{E} \left[\int_0^\infty e^{-\beta t} U(c_t) dt \right],$$

where $U: (0, \infty) \mapsto \mathbb{R}$ is a utility function. We assume that U is C^2 , increasing, strictly concave, and we denote its convex conjugate by,

$$\tilde{U}(\tilde{c}) := \sup_{c>0} \{U(c) - c\tilde{c}\}, \quad \tilde{c} \in \mathbb{R}.$$

2.2. Dynamic Programming Equation

The dynamic programming equation corresponding to the singular stochastic control problem v^ϵ involves the following differential operators. Let:

$$\mathcal{L} := \mu \cdot (\mathbf{D}_s + \mathbf{D}_y) + r\mathbf{D}_x + \frac{1}{2} \text{Tr} [\sigma \sigma^\top (\mathbf{D}_{yy} + \mathbf{D}_{ss} + 2\mathbf{D}_{sy})], \quad (2.1)$$

and for $i, j = 1, \dots, d$,

$$\begin{aligned} \mathbf{D}_x &:= x \frac{\partial}{\partial x}, & \mathbf{D}_s^i &:= s^i \frac{\partial}{\partial s^i}, & \mathbf{D}_y^i &:= y^i \frac{\partial}{\partial y^i}, \\ \mathbf{D}_{ss}^{i,j} &:= s^i s^j \frac{\partial^2}{\partial s^i \partial s^j}, & \mathbf{D}_{yy}^{i,j} &:= y^i y^j \frac{\partial^2}{\partial y^i \partial y^j}, & \mathbf{D}_{sy}^{i,j} &:= s^i y^j \frac{\partial^2}{\partial s^i \partial y^j}, \end{aligned}$$

$\mathbf{D}_s = (\mathbf{D}_s^i)_{1 \leq i \leq d}$, $\mathbf{D}_y = (\mathbf{D}_y^i)_{1 \leq i \leq d}$, $\mathbf{D}_{yy} := (\mathbf{D}_{yy}^{i,j})_{1 \leq i, j \leq d}$, $\mathbf{D}_{ss} := (\mathbf{D}_{ss}^{i,j})_{1 \leq i, j \leq d}$, $\mathbf{D}_{sy} := (\mathbf{D}_{sy}^{i,j})_{1 \leq i, j \leq d}$. Moreover, for a smooth scalar function $(s, x, y) \in \mathbb{R}_+^d \times \mathbb{R} \times \mathbb{R}^d \mapsto \varphi(x, y)$, we set

$$\varphi_x := \frac{\partial \varphi}{\partial x} \in \mathbb{R}, \quad \varphi_y := \frac{\partial \varphi}{\partial y} \in \mathbb{R}^d.$$

Theorem 2.1. *Assume that the value function v^ϵ is locally bounded. Then, v^ϵ is a viscosity solution of the dynamic programming equation in $\mathbb{R}_+^d \times K_\epsilon$,*

$$\min_{(i,j) \in \mathcal{J}} \left\{ \beta v^\epsilon - \mathcal{L}v^\epsilon - \tilde{U}(v_x^\epsilon), \Lambda_{i,j}^\epsilon \cdot (v_x^\epsilon, v_y^\epsilon) \right\} = 0, \quad \Lambda_{i,j}^\epsilon := e_i - e_j + \epsilon^3 \lambda^{i,j} e_i, \quad (i, j) \in \mathcal{J}. \quad (2.2)$$

Moreover, v^ϵ is concave in (x, y) and converges to the Merton value function $v := v^0$, as $\epsilon > 0$ tends to zero.

The Merton value function $v = v^0$ corresponds to the limiting case $\epsilon = 0$ where the transfers between assets are not subject to transaction costs. Our subsequent analysis assumes that v is smooth, which can be verified under slight conditions on the coefficients. We refer to [40] for the well-known properties and related equations satisfied by v . For future reference, let us however denote by \mathbf{y} and \mathbf{c} the optimal controls in the Merton problem which satisfy

$$\mathbf{c}(s, z) := -\tilde{U}'_1(v_z(s, z)) = (U'_1)^{-1}(v_z(s, z)) \quad (2.3)$$

$$-v_{zz}(s, z) \sigma \sigma^T(s) \mathbf{y}(s, z) := (\mu - r \mathbf{1}_d)(s) v_z(s, z) + \sigma \sigma^T(s) \mathbf{D}_{sz} v(s, z), \quad (2.4)$$

where we used the change of variables $z := x + y \cdot \mathbf{1}_d$.

2.3. Formal Asymptotics

Let us define the following change of variable for any $(x, y) \in \mathbb{R}^{d+1}$

$$\xi_i := \zeta_i^\epsilon(s, z) = \frac{y_i - \mathbf{y}_i(s, z)}{\epsilon}.$$

Here, we recall the formal first order expansion derived in [40]:

$$v^\epsilon(s, x, y) = v(s, z) - \epsilon^2 u(s, z) - \epsilon^4 w(s, z, \xi) + o(\epsilon^2), \quad (2.5)$$

where u is solution of the second corrector equation:

$$\mathcal{A}u := \beta u - \mathcal{L}^0 u - (rz + \mathbf{y} \cdot (\mu - r \mathbf{1}_d) - \mathbf{c}) u_z - \frac{1}{2} |\sigma^T \mathbf{y}|^2 u_{zz} - \sigma \sigma^T \mathbf{y} \cdot \mathbf{D}_{sz} u = a,$$

and where (w, a) is the solution of the first corrector equation:

$$\max_{(i,j) \in \mathcal{F}} \max \left\{ \frac{|\sigma^T(s)\xi|^2}{2} v_{zz}(s, z) - \frac{1}{2} \text{Tr} [\alpha \alpha^T(s, z) w_{\xi\xi}(s, z, \xi)] + a(s, z); \right. \\ \left. -\lambda^{i,j} v_z(s, z) + \frac{\partial w}{\partial \xi_i}(s, z, \xi) - \frac{\partial w}{\partial \xi_j}(s, z, \xi) \right\} = 0,$$

where $\xi \in \mathbb{R}^d$ is the dependent variable, while $(s, z) \in (0, \infty)^d \times \mathbb{R}^+$ are fixed, and the diffusion coefficient is given by

$$\alpha(s, z) := [(I_d - \mathbf{y}_z(s, z) \mathbf{1}_d^T) \text{diag}[\mathbf{y}(s, z)] - \mathbf{y}_s^T(s, z) \text{diag}[s]] \sigma(s).$$

The expansion 2.5 was proved rigorously in [40] in the one-dimensional case, with a crucial use of the explicit solution of the first corrector equation in one space dimension. Our objective in this paper is to show that the above expansion is valid in the present d -dimensional framework where no explicit solution of the first corrector equation is available anymore.

We finally recall the from [40] the following normalization. Set

$$\eta(s, z) := -\frac{v_z(s, z)}{v_{zz}(s, z)}, \quad \rho := \frac{\xi}{\eta(s, z)}, \quad \bar{w}(s, z, \rho) := \frac{w(s, z, \eta(s, z)\rho)}{\eta(s, z)v_z(s, z)}, \\ \bar{a}(s, z) := \frac{a(s, z)}{\eta(s, z)v_z(s, z)}, \quad \bar{\alpha}(s, z) := \frac{\alpha(s, z)}{\eta(s, z)},$$

so that the corrector equations with variable $\rho \in \mathbb{R}^d$ have the form,

$$\max_{(i,j) \in \mathcal{F}} \max \left\{ \frac{|\sigma^T(s)\rho|^2}{2} - \frac{1}{2} \text{Tr} [\bar{\alpha} \bar{\alpha}^T(s, z) \bar{w}_{\rho\rho}(s, z, \rho)] + \bar{a}(s, z); \right. \\ \left. -\lambda^{i,j} + \frac{\partial \bar{w}}{\partial \rho_i}(s, z, \rho) - \frac{\partial \bar{w}}{\partial \rho_j}(s, z, \rho) \right\} = 0 \quad (2.6)$$

$$\mathcal{A}u(s, z) = v_z(s, z)\eta(s, z)\bar{a}(s, z). \quad (2.7)$$

Notice that we will always use the normalization $\bar{w}(s, z, 0) = 0$.

3. The Main Results

3.1. The First Corrector Equation

We first state the existence and uniqueness of a solution to the first corrector equation (2.6), within a convenient class of functions. We also state the main properties of the solution which will be used later. We will often make use of the language of ergodic control theory, and say that \bar{w} is a solution of (2.6) with eigenvalue \bar{a} .

Consider the following closed convex subset of \mathbb{R}^d , and the corresponding support function

$$C := \{\rho \in \mathbb{R}^d : -\lambda^{j,i} \leq \rho_i - \rho_j \leq \lambda^{i,j}, (i, j) \in \mathcal{F}\}, \quad \delta_C(\rho) := \sup_{u \in C} u \cdot \rho, \quad \rho \in \mathbb{R}^d,$$

with the convention that $\rho_0 = 0$. Then, C is a bounded convex polyhedral containing 0, which corresponds to the intersection of $d(d+1)$ hyperplanes. Furthermore, the gradient constraints of the corrector equation (2.6) is exactly that $D\bar{w} \in C$. In this respect, (2.6) is closely related to the variational inequality studied by Menaldi et al. [28] and Hynd [21], where the gradient is restricted to lie in the closed unit ball of \mathbb{R}^d (we also refer to [22] for related variational inequalities with general gradient constraints).

The wellposedness of the first corrector equation is stated in the following two results. Since the variables (s, z) are frozen in the equation (2.6), we omit the dependence on them.

Theorem 3.1 (First corrector equation: comparison). *Assume that $\bar{\alpha}$ is non-degenerate. Suppose w_1 is a viscosity subsolution of (2.6) with eigenvalue a_1 and that w_2 is a viscosity supersolution of (2.6) with eigenvalue a_2 . Assume further that*

$$\overline{\lim}_{|\rho| \rightarrow +\infty} \frac{w_1(\rho)}{\delta_C(\rho)} \leq 1 \leq \underline{\lim}_{|\rho| \rightarrow +\infty} \frac{w_2(\rho)}{\delta_C(\rho)}.$$

Then, $a_1 \leq a_2$.

The proof of this result is given in Section 5.1.

Theorem 3.2 (First corrector equation: existence). *Assume that $\bar{\alpha}$ is non-degenerate. Then, there exists a solution $\bar{w} \in C^{1,1}$ of the equation (2.6) with eigenvalue $\bar{a} \in \mathbb{R}_+$, with $\bar{w}(0) = 0$ and satisfying the growth condition $\lim_{|\rho| \rightarrow \infty} (\bar{w}/\delta_C)(\rho) = 1$. Moreover,*

- \bar{w} is convex.
- The set $\mathcal{O}_0 := \{\rho \in \mathbb{R}^d, D\bar{w}(\rho) \in \text{int}(C)\}$ is open and bounded, $\bar{w} \in C^\infty(\mathcal{O}_0)$ and

$$\bar{w}(\rho) = \inf_{y \in \mathcal{O}_0} \{\bar{w}(y) + \delta_C(\rho - y)\}, \quad \text{for all } \rho \in \mathbb{R}^d.$$

- $\bar{w}(\rho)$ attains its minimum in ρ at some point ρ^* in \mathcal{O}_0 .
- There is a constant $M > 0$ such that $0 \leq D^2\bar{w}(\rho) \leq M\mathbf{1}_{\bar{\mathcal{O}}_0}(\rho)$ for a.e. $\rho \in \mathbb{R}^d$.

The proof of this result will be reported in Section 5.2. In the one-dimensional context $d = 1$, the unique solution of the first corrector equation is easily obtained in explicit form in [40]. For higher dimension $d \geq 2$, in general, no such explicit expression is available anymore. The following example illustrates a particular structure of the parameters of the problem which allows to obtain an explicit solution.

Example 3.1. Suppose the equation is of the form,

$$\max_{0 \leq i \leq d} \max \left\{ -\frac{c_1^* |\rho|^2}{2} - \frac{c_2^*}{2} \Delta \hat{w}(\rho) + \hat{a}, -\hat{\lambda}^i + \frac{\partial \hat{w}}{\partial \rho_i}(\rho), -\tilde{\lambda}^i - \frac{\partial \hat{w}}{\partial \rho_i}(\rho) \right\} = 0, \quad (3.1)$$

with given positive constants $c_1^*, c_2^*, \hat{\lambda}^i, \tilde{\lambda}^i$ and the normalization $\hat{w}(0) = 0$. This corresponds to the case when σ and $\bar{\alpha}$ are multiples of the identity matrix and we are only allowed to make transactions to and from the cash account, i.e., when $\lambda^{i,j} = \infty$

as soon as i and j are both different from 0. Then, the unique solution \hat{w} is given as,

$$\hat{w}(\rho) = \sum_{i=1}^d \tilde{w}_i(\rho_i),$$

where \tilde{w}_i is the explicit solution of the one dimensional problem constructed in [40]. Moreover, \hat{a} is an explicit constant independent of z .

Notice that for the corrector equations, σ and $\bar{\alpha}$ cannot be specified independent of each other. However, the above explicit solution will be used as an upper bound for the unique solution of the corrector equation.

3.2. Assumptions

This section collects all assumptions which are needed for our main results, and comments on them. In particular, they are verified in one space dimension in [40]. In subsection 3.2.4 below, we show that they also hold in the multi-dimensional power utility case.

3.2.1. Assumptions on the Merton Value Function.

Assumption 3.1 (Smoothness). v and \mathbf{y} are C^2 in $(0, \infty)^{d+1}$, $v_z > 0$, $\mathbf{y}_z^i > 0$, $i \leq d$, on $(0, \infty)^{d+1}$, and there exist $c_0, c_1 > 0$ such that

$$c_0 \leq \mathbf{y}_z \cdot \mathbf{1}_d \leq 1 - c_0 \quad \text{and} \quad \alpha \alpha^T \geq c_1 I_d, \quad \text{on} \quad (0, \infty)^{d+1}.$$

Notice that this assumption is verified in the case of the Black-Scholes model with power utility.

3.2.2. Local Boundedness. As in [40], we define

$$\bar{u}^\epsilon(s, x, y) := \frac{v(s, z) - v^\epsilon(s, x, y)}{\epsilon^2}, \quad s \in \mathbb{R}_+^d, \quad (x, y) \in K_\epsilon. \quad (3.2)$$

Following the classical approach of Barles and Perthame, we introduce the relaxed semi-limits

$$u^*(s, x, y) := \overline{\lim}_{(\epsilon, s', x', y') \rightarrow (0, s, x, y)} \bar{u}^\epsilon(s', x', y'), \quad u_*(s, x, y) := \underline{\lim}_{(\epsilon, s', x', y') \rightarrow (0, s, x, y)} \bar{u}^\epsilon(s', x', y').$$

The following assumption is verified in Lemma 3.1 below, in the power utility context with constant coefficients.

Assumption 3.2 (Local bound). *The family of functions \bar{u}^ϵ is locally uniformly bounded from above.*

This assumption states that for any $(s_0, x_0, y_0) \in (0, \infty)^d \times \mathbb{R} \times \mathbb{R}^d$ with $x_0 + y_0 \cdot \mathbf{1}_d > 0$, there exist $r_0 = r_0(s_0, x_0, y_0) > 0$ and $\epsilon_0 = \epsilon_0(s_0, x_0, y_0) > 0$ so that

$$b(s_0, x_0, y_0) := \sup\{u^\epsilon(s, x, y) : (s, x, y) \in B_{r_0}(s_0, x_0, y_0), \epsilon \in (0, \epsilon_0]\} < \infty, \quad (3.3)$$

where $B_{r_0}(s_0, x_0, y_0)$ denotes the open ball with radius r_0 , centered at (s_0, x_0, y_0) .

3.2.3. Assumptions on the Corrector Equations. Let b be as in 3.3, and set

$$B(s, z) := b(s, z - \mathbf{y}(s, z) \cdot \mathbf{1}_d, \mathbf{y}(s, z)), \quad s \in (0, \infty)^d, \quad z \geq 0. \quad (3.4)$$

Assumption 3.3 (Second corrector equation: comparison). *For any upper-semicontinuous (resp. lower-semicontinuous) viscosity subsolution (resp. supersolution) u_1 (resp. u_2) of (2.7) in $(0, \infty)^{d+1}$ satisfying the growth condition $|u_i| \leq B$ on $(0, \infty)^{d+1}$, $i = 1, 2$, we have $u_1 \leq u_2$ in $(0, \infty)^{d+1}$.*

In the above comparison, notice that the growth of the supersolution and the subsolution is controlled by the function B which is defined in 3.4 by means of the local bound function b . In particular, B controls the growth both at infinity and near the origin. Also note that in the power case, by homothety, there are constants $0 < a_* \leq a^* < \infty$ so that $u_*(z) = a_* z^p$, $u^*(z) = a^* z^p$ where p is the power of the utility function. Then, the comparison follows from a direct verification argument. In the general case, \mathcal{A} is the infinitesimal generator of the optimal wealth process in the Merton problem. Hence, one may also use the probabilistic arguments combined with the known estimates of the optimal wealth process to prove comparison.

Our next assumption concerns the continuity of the solution (w, a) of the first corrector equation in the parameters (s, z) . Recall that $w = \eta v_z \bar{w}$ and $a = \eta v_z \bar{a}$.

Assumption 3.4 (First corrector equation: regular dependence on the parameters). *The set $\mathcal{C}_0(s, z)$ and the functions $a(s, z)$ and $\rho^*(s, z)$ are continuous in (s, z) . Moreover, both w and $\tilde{w}(\cdot, \xi) := w(\cdot, \xi) - w(\cdot, \eta(\cdot)\rho^*(\cdot))$ are C^2 in (s, z) , and satisfy the following estimates*

$$(|\varpi_s| + |\varpi_{ss}| + |\varpi_z| + |\varpi_{sz}| + |\varpi_{zz}|)(s, z, \xi) \leq C(s, z)(1 + |\xi|) \quad (3.5)$$

$$(|\varpi_\xi| + |\varpi_{s\xi}| + |\varpi_z \xi|)(s, z, \xi) \leq C(s, z), \quad (3.6)$$

for $\varpi = w$ or \tilde{w} and where $C(s, z)$ is a continuous function depending on the Merton value function and its derivatives.

Notice that by the stability of viscosity solutions, the function w is clearly continuous in (s, z) . Moreover, Assumption 3.4 is satisfied when we consider the constant coefficients and the power utility function. Indeed, in that case, there is no dependence in the s variable, as emphasized in Lemma 3.1, and the dependence in z can be factored out by homogeneity.

3.2.4. Power Utility. In this subsection, we verify the above assumptions for the power utility with Black-Scholes dynamics. We also observe that this is just one possible set of assumptions and the below arguments can be modified to obtain the same result under other conditions.

Lemma 3.1. *Suppose that U is a power utility and the stock dynamics is Black-Scholes, i.e., the functions r, μ, σ are constants. Then, all above assumptions hold.*

Proof. Since the value function is independent of the s variable and positively homogenous in the z variable, Assumptions 3.1, 3.3, 3.4 are directly verified. We continue by proving Assumption 3.2.

Let \hat{v}^ϵ be the value function of the same utility maximization problem with transactions are forced to go through the cash account, i.e., $\lambda^{i,j} = \infty$ for all $1 \leq i \neq j \leq d$. Set $\hat{u}^\epsilon := (v - \hat{v}^\epsilon)/\epsilon^2$. Then, clearly $\hat{v}^\epsilon \leq v^\epsilon$ and $0 \leq u^\epsilon \leq \hat{u}^\epsilon$. Therefore, without loss of generality, we may assume that $\lambda^{i,j} = \infty$ for all $1 \leq i \neq j \leq d$. Recall that we always assume that $0 < \lambda^{i,0}, \lambda^{0,i} < \infty$ for all $1 \leq i \leq d$.

The homotheticity of the utility function implies that both the Merton value function v and the solution u of the second corrector equation are homothetic as well. For a large positive constant K to be chosen later, set

$$V^{\epsilon,K}(z, \xi) := v(z) - \epsilon^2 K u(z) - \epsilon^4 W(z, \xi),$$

where $W(z, \xi) := z v'(z) \hat{w}(\xi/z)$, and \hat{w} solves (3.1) with constants chosen so that

$$c_1^* I_d \geq \sigma \sigma^T, \quad c_2^* I_d \geq \bar{\alpha} \bar{\alpha}^T, \quad \hat{\lambda}^i = \tilde{\lambda}^i = 2\bar{\lambda} := \max_{(i,j) \in \mathcal{J}} \lambda^{i,j}.$$

Note that W is explicit and is twice continuously differentiable. We continue by showing that for large K , $V^{\epsilon,K}$ is a subsolution of the dynamic programming equation (2.2), which would imply that $V^{\epsilon,K} \leq v^\epsilon$ by the comparison result for the equation (2.2) (see [24] Theorem 4.3.2, Proposition 4.3.4 and the subsequent discussion). A straightforward calculation, as in the proof of Lemma 7.2 in [40], shows that the gradient constraints

$$\Lambda_{i,0}^\epsilon \cdot (V_x^{\epsilon,K}, V_{y_i}^{\epsilon,K}) \leq 0, \quad \text{holds whenever} \quad 2\bar{\lambda} + \frac{\partial \hat{w}}{\partial \rho_i}(\rho) \leq 0.$$

Similarly,

$$\Lambda_{0,i}^\epsilon \cdot (V_x^{\epsilon,K}, V_{y_i}^{\epsilon,K}) \leq 0, \quad \text{holds whenever} \quad -2\bar{\lambda} + \frac{\partial \hat{w}}{\partial \rho_i}(\rho) \leq 0.$$

Assume therefore that the elliptic part in the equation (3.1) holds. We claim that for a large constant K ,

$$\mathcal{J}(V^{\epsilon,K}) := \beta V^{\epsilon,K} - \mathcal{L}V^{\epsilon,K} - \tilde{U}(V_x^{\epsilon,K}) \leq 0.$$

We proceed exactly as in subsection 4.2 below to arrive at

$$\mathcal{J}(V^{\epsilon,K}) = \epsilon^2 \left[-\frac{1}{2} |\sigma(s)\xi| v_{zz} + \frac{1}{2} \text{Tr} [\alpha \alpha^T(s, z) W_{\xi\xi}(s, z, \xi)] - K \mathcal{A}u(z) + \mathcal{R}^\epsilon(z, \xi) \right].$$

It is clear that $|\mathcal{R}^\epsilon(z, \xi)| \leq k^* \epsilon z v'(z)$ for some constant k^* . We now use the elliptic part of the equation (3.1) and the choices of c_i^* to conclude that in this region,

$$-\frac{1}{2} |\sigma^T(s)\xi| v_{zz} + \frac{1}{2} \text{Tr} [\alpha \alpha^T(s, z) W_{\xi\xi}(s, z, \xi)] \leq z v'(z) \hat{a}.$$

Also by homotheticity, $a(z) = zv'(z)a_0$ for some $a_0 > 0$. Hence,

$$\mathcal{F}(V^{\epsilon,K}) \leq \epsilon^2 zv'(z) [\hat{a} - Ka_0 + \epsilon k^*] \leq 0,$$

provided that K is sufficiently large. Since $V^{\epsilon,K}$ is a smooth subsolution of the dynamic programming equation, we conclude by using the standard verification argument. \square

3.3. The First Order Expansion Result

The main result of this paper is the following d -dimensional extension of [40].

Theorem 3.3. *Under Assumptions 3.1, 3.2, 3.3, and 3.4, the sequence $\{\bar{u}^\epsilon\}_{\epsilon>0}$, defined in (3.2), converges locally uniformly to the function u defined in (2.7).*

Proof. In Section 4, we will show that, the semi-limits u_* and u^* are viscosity supersolution and subsolution, respectively, of (2.7). Then, by the comparison Assumption 3.3, we conclude that $u^* \leq u \leq u_*$. Since the opposite inequality is obvious, this implies that $u^* = u_* = u$. The local uniform convergence follows immediately from this and the definitions. \square

4. Convergence

In the rest of the paper, we denote for any function $f(s, x, y)$:

$$\hat{f}(s, z, \zeta) := f(s, z - \mathbf{y}(s, z) \cdot \mathbf{1}_d, \epsilon \zeta + \mathbf{y}(s, z)).$$

This section is dedicated to the proof of our main result, Theorem 3.3. Let:

$$u^\epsilon(s, x, y) := \bar{u}^\epsilon(s, x, y) - \epsilon^2 w(s, z, \zeta), \quad s \in \mathbb{R}_+^d, \quad (x, y) \in K_\epsilon.$$

4.1. First Estimates and Properties

We start by obtaining several estimates of u^ϵ . Set

$$\bar{\lambda} := \max_{(i,j) \in \mathcal{J}} \lambda^{i,j}, \quad \underline{\lambda} := \min_{(i,j) \in \mathcal{J}} \lambda^{i,j}.$$

We also recall that L is the upper bound of the set C . The following two Lemmas can be proved exactly as Lemmas 6.1 and 6.2 in [40]. The multidimensional setting only renders one estimate more difficult to obtain, so that we will report only its proof in the appendix.

Lemma 4.1. *For $(\epsilon, s, x, y) \in (0, +\infty) \times (0, +\infty)^d \times K_\epsilon$, and $z := x + y \cdot \mathbf{1}_d$, we have*

$$u^\epsilon(s, x, y) \geq -\epsilon Lv_z(s, z) |y - \mathbf{y}(s, z)|.$$

Consequently, under Assumption 3.2, we have for all $(\epsilon, s, x, y) \in (0, +\infty) \times (0, +\infty)^d \times K_\epsilon$

$$0 \leq u_*(s, x, y) \leq u^*(s, x, y) < +\infty.$$

The next Lemma proves that the relaxed semi-limits are only functions of (s, z) .

Lemma 4.2. *Let Assumptions 3.1 and 3.2 hold. Then, u^* and u_* are only functions of (s, z) . Furthermore, we have*

$$\begin{aligned} u_*(s, z) &= \lim_{(\epsilon, s', z') \rightarrow (0, s, z)} u^\epsilon(s', z' - \mathbf{y}(s', z') \cdot \mathbf{1}_d, \mathbf{y}(s', z')) \\ u^*(s, z) &= \overline{\lim}_{(\epsilon, s', z') \rightarrow (0, s, z)} u^\epsilon(s', z' - \mathbf{y}(s', z') \cdot \mathbf{1}_d, \mathbf{y}(s', z')). \end{aligned}$$

4.2. Remainder Estimate

In this section, we isolate an estimate which will be needed at various occasions in the subsequent proofs. The following calculation extends the estimate of Section 4.2 in [40]. For any function

$$\Psi^\epsilon(s, x, y) := v(s, z) - \epsilon^2 \phi(s, z) - \epsilon^4 \varpi(s, z, \xi),$$

with smooth ϕ and v such that v also verifies the estimates 3.5, we have

$$\begin{aligned} \mathcal{J}(\Psi^\epsilon)(s, x, y) &:= \left(\beta \Psi^\epsilon - \mathcal{L} \Psi^\epsilon - \tilde{U}(\Psi_x^\epsilon) \right) (s, x, y) \\ &= \epsilon^2 \left[-\frac{1}{2} |\sigma(s) \xi| v_{zz} + \frac{1}{2} \text{Tr} [\alpha \alpha^T (s, z) \varpi_{\xi\xi}(s, z, \xi)] \right. \\ &\quad \left. - \mathcal{A} \phi(s, z) + \mathcal{R}^\epsilon(s, z, \xi) \right]. \end{aligned}$$

Similarly as in [40], direct but tedious calculations provides the following estimate:

$$\begin{aligned} |\mathcal{R}^\epsilon(s, x, y)| &\leq \epsilon \left(|\mu - r \cdot \mathbf{1}_d| |\xi| |\phi_z| + \frac{|\sigma|^2}{2} (2|\mathbf{y}| |\xi| + \epsilon \xi^2) |\phi_{zz}| + |\sigma|^2 |\xi| |s| |\phi_{sz}| \right) (s, z) \\ &\quad + \epsilon C(s, z) \left(1 + \epsilon |\xi| + \epsilon^2 |\xi|^2 + \epsilon^3 |\xi|^3 \right) \\ &\quad + \epsilon^{-2} \left| \tilde{U}(\Psi_x^\epsilon) - \tilde{U}(v_z) - (\Psi_x^\epsilon - v_z) \tilde{U}'(v_z) \right|, \end{aligned}$$

for some continuous function $C(s, z)$. Now using the fact \tilde{U} is C^1 and convex and the estimates assumed for v , we obtain

$$\begin{aligned} |\mathcal{R}^\epsilon(s, x, y)| &\leq \epsilon \left(|\mu - r \cdot \mathbf{1}_d| |\xi| |\phi_z| + \frac{|\sigma|^2}{2} (2|\mathbf{y}| |\xi| + \epsilon \xi^2) |\phi_{zz}| + |\sigma|^2 |\xi| |s| |\phi_{sz}| \right) (s, z) \\ &\quad + \epsilon C(s, z) \left(1 + \epsilon |\xi| + \epsilon^2 |\xi|^2 + \epsilon^3 |\xi|^3 \right) \\ &\quad + \epsilon^2 (|\phi_z| + \epsilon C(s, z) (1 + \epsilon |\xi|))^2 \tilde{U}''(v_z + \epsilon^2 |\phi_z| + \epsilon^3 C(s, z) (1 + \epsilon |\xi|)). \end{aligned}$$

4.3. Viscosity Subsolution Property

In this Section, we prove

Proposition 4.1. *Under Assumptions 3.1, 3.2 and 3.4, the function u^* is a viscosity subsolution of the second corrector equation.*

Proof. Let $(s_0, z_0, \phi) \in (0, +\infty)^d \times (0, +\infty) \times C^2((0, +\infty)^d \times (0, +\infty))$ be such that

$$(u^* - \phi)(s_0, z_0) > (u^* - \phi)(s, z), \quad \text{for all } (s, z) \in (0, +\infty)^d \times (0, +\infty) \setminus \{(s_0, z_0)\}. \quad (4.1)$$

By definition of viscosity subsolutions, we need to show that

$$\mathcal{A}\phi(s_0, z_0) - a(s_0, z_0) \leq 0.$$

We will proceed in several steps.

Step 1: First of all, we know from Lemma 4.2 that there exists a sequence (s^ϵ, z^ϵ) which realizes the lim sup for \hat{u}^ϵ , that is to say

$$(s^\epsilon, z^\epsilon) \xrightarrow{\epsilon \downarrow 0} (s_0, z_0) \quad \text{and} \quad \hat{u}^\epsilon(s^\epsilon, z^\epsilon, 0) \xrightarrow{\epsilon \downarrow 0} u^*(s_0, z_0).$$

It follows then easily that $l_*^\epsilon := \hat{u}^\epsilon(s^\epsilon, z^\epsilon, 0) - \phi(s^\epsilon, z^\epsilon) \xrightarrow{\epsilon \downarrow 0} 0$, and

$$(x^\epsilon, y^\epsilon) = (z^\epsilon - \mathbf{y}(s^\epsilon, z^\epsilon) \cdot \mathbf{1}_d, \mathbf{y}(s^\epsilon, z^\epsilon)) \xrightarrow{\epsilon \downarrow 0} (x_0, y_0) := (z_0 - \mathbf{y}(s_0, z_0) \cdot \mathbf{1}_d, \mathbf{y}(s_0, z_0)).$$

Now recall from Assumption 3.2 that u^ϵ is locally bounded from above. This implies the existence of $r_0 := r_0(s_0, x_0, y_0) > 0$ and $\epsilon_0 := \epsilon_0(s_0, x_0, y_0) > 0$ verifying

$$b_* := \sup \{u^\epsilon(s, x, y), (s, x, y) \in B_0, \epsilon \in (0, \epsilon_0]\} < +\infty, \quad (4.2)$$

where $B_0 := B_{r_0}(s_0, x_0, y_0)$ is the open ball with radius r_0 and center (s_0, x_0, y_0) . Moreover, notice that we can always decrease r_0 so that $r_0 \leq z_0/2$, which then implies that B_0 does not cross the line $z = 0$. Now for any $(\epsilon, \delta) \in (0, 1]^2$, we define $\Psi^{\epsilon, \delta}$ and the corresponding $\hat{\Psi}^{\epsilon, \delta}$ by

$$\hat{\Psi}^{\epsilon, \delta}(s, z, \xi) := v(s, z) - \epsilon^2 \left(l_*^\epsilon + \phi(s, z) + \hat{\Phi}^\epsilon(s, z, \xi) \right) - \epsilon^4 (1 + \delta) \tilde{w}(s, z, \xi),$$

where the function $\hat{\Phi}^\epsilon$ and the corresponding Φ^ϵ are given by:

$$\hat{\Phi}^\epsilon(s, x, y) := c_0 \left((s - s^\epsilon)^4 + (z - z^\epsilon)^4 + \epsilon^4 \tilde{w}^4(s, z, \xi) \right),$$

and $c_0 > 0$ is a constant chosen large enough in order to have for ϵ small enough

$$\Phi^\epsilon \geq 1 + b_* - \phi, \quad \text{on } B_0 \setminus B_1, \quad \text{where } B_1 := B_{\frac{r_0}{2}}(s_0, x_0, y_0). \quad (4.3)$$

We emphasize that the constant c_0 may depend on $(\phi, s_0, x_0, y_0, \delta)$ but not on ϵ , and that *a priori* the function $\hat{\Psi}^{\epsilon, \delta}$ is not C^2 in ξ , because the function w is only in $C^{1,1}$. This is a major difference with the one-dimension case treated in [40].

Step 2: We now prove that for all sufficiently small ϵ and δ , the difference $(v^\epsilon - \Psi^{\epsilon,\delta})$ has a local minimizer in B_0 . First, notice that this is equivalent to showing that the following quantity has a local minimizer in B_0

$$\begin{aligned} I^{\epsilon,\delta}(s, x, y) &:= \frac{v^\epsilon(s, x, y) - \Psi^{\epsilon,\delta}(s, x, y)}{\epsilon^2} \\ &= -u^\epsilon(s, x, y) + l_*^\epsilon + \phi(s, z) + \Phi^\epsilon(s, x, y) + \epsilon^2 \delta w(s, z, \zeta) \\ &\quad - \epsilon^2 w(s, z, \eta(s, z) \rho^*(s, z)). \end{aligned}$$

By (4.3), the fact that $\tilde{w} \geq 0$ and by continuity of η and ρ^* , we have for any $(s, x, y) \in \partial B_0$

$$I^{\epsilon,\delta}(s, x, y) \geq -u^\epsilon(s, x, y) + l_*^\epsilon + 1 + b_* - \epsilon^2 w(s, z, \eta(s, z) \rho^*(s, z)) \geq \frac{1}{2} + l_*^\epsilon > 0,$$

for ϵ small enough. Moreover, since $I^{\epsilon,\delta}(s^\epsilon, x^\epsilon, y^\epsilon) \rightarrow 0$ as ϵ goes to 0, this implies that $I^{\epsilon,\delta}$ has a local minimizer $(\tilde{s}^\epsilon, \tilde{x}^\epsilon, \tilde{y}^\epsilon)$ in B_0 , and we introduce the corresponding

$$\tilde{z}^\epsilon := \tilde{x}^\epsilon + \tilde{y}^\epsilon \cdot \mathbf{1}_d, \quad \text{and} \quad \tilde{\zeta}^\epsilon := \frac{\tilde{y}^\epsilon - \mathbf{y}(\tilde{s}^\epsilon, \tilde{z}^\epsilon)}{\epsilon}.$$

We then have

$$\begin{aligned} \min_{(s, z, \zeta)} (\hat{v}^\epsilon - \hat{\Psi}^{\epsilon,\delta})(s, z, \zeta) &= (\hat{v}^\epsilon - \Psi^{\epsilon,\delta})(\tilde{s}^\epsilon, \tilde{z}^\epsilon, \tilde{\zeta}^\epsilon) \leq 0, \quad |\tilde{s}^\epsilon - s_0| + |\tilde{z}^\epsilon - z_0| \\ &\leq r_0, \quad \left| \frac{\tilde{\zeta}^\epsilon}{\epsilon} \right| \leq \frac{r_1}{\epsilon}, \end{aligned}$$

for some constant r_1 . We now use the viscosity supersolution property of v^ϵ . Since $\Psi^{\epsilon,\delta}$ is C^1 , we obtain from the first order operator in the dynamic programming equation that:

$$\Lambda_{i,j}^\epsilon \cdot (\Psi_x^{\epsilon,\delta}, \Psi_y^{\epsilon,\delta})(\tilde{s}^\epsilon, \tilde{x}^\epsilon, \tilde{y}^\epsilon) \geq 0 \quad \text{for all } (i, j) \in \mathcal{J}. \quad (4.4)$$

Step 3: In this step, we show that for ϵ small enough, we have

$$\tilde{\rho}^\epsilon := \frac{\tilde{\zeta}^\epsilon}{\eta(\tilde{s}^\epsilon, \tilde{z}^\epsilon)} \in \mathcal{O}_0(\tilde{s}^\epsilon, \tilde{z}^\epsilon). \quad (4.5)$$

where $\mathcal{O}_0(s, z)$ is the open set of Proposition 3.2. We argue by contradiction assuming that there exists some sequence $\epsilon_n \rightarrow 0$ such that $\tilde{\rho}^{\epsilon_n} \notin \mathcal{O}_0(\tilde{s}^{\epsilon_n}, \tilde{z}^{\epsilon_n})$. This implies that

$$-\lambda_{i_0, j_0}^{\epsilon_n} + (\partial_{i_0}^{\epsilon_n} \bar{w} - \partial_{j_0}^{\epsilon_n} \bar{w})(\tilde{s}^{\epsilon_n}, \tilde{z}^{\epsilon_n}, \tilde{\rho}^{\epsilon_n}) = 0 \quad \text{for some } (i_0, j_0) \in \mathcal{J}. \quad (4.6)$$

By the gradient constraints (4.4), and the boundedness of $(\tilde{s}^{\epsilon_n}, \tilde{z}^{\epsilon_n}, \epsilon_n \tilde{\xi}^{\epsilon_n})_n$, we directly compute that:

$$\begin{aligned} & -4C\epsilon_n^2(\epsilon_n \tilde{w})^3(\tilde{s}^{\epsilon_n}, \tilde{z}^{\epsilon_n}, \tilde{\xi}^{\epsilon_n}) \left(w_{\xi_0^n} - w_{\tilde{\xi}_0^n} \right) (\tilde{s}^{\epsilon_n}, \tilde{z}^{\epsilon_n}, \tilde{\xi}^{\epsilon_n}) \\ & + \epsilon_n^3 v_z(\tilde{s}^{\epsilon_n}, \tilde{z}^{\epsilon_n}) \left[\lambda_{j_0^n}^{\tilde{j}_0^n} - (1 + \delta)(\partial_{j_0^n} \bar{w} - \partial_{\tilde{j}_0^n} \bar{w})(\tilde{s}^{\epsilon_n}, \tilde{z}^{\epsilon_n}, \tilde{\rho}^{\epsilon_n}) \right] + o(\epsilon_n^3) \geq 0. \end{aligned} \quad (4.7)$$

Using (4.6) and the non-negativity of \tilde{w} , this implies

$$\begin{aligned} 0 & \leq -4c_0 \lambda_{j_0^n}^{\tilde{j}_0^n} \epsilon_n^2 (\epsilon_n \tilde{w})^3 (\tilde{s}^{\epsilon_n}, \tilde{z}^{\epsilon_n}, \tilde{\xi}^{\epsilon_n}) - \delta \lambda_{j_0^n}^{\tilde{j}_0^n} \epsilon_n^3 v_z(\tilde{s}^{\epsilon_n}, \tilde{z}^{\epsilon_n}) + o(\epsilon_n^3) \\ & \leq -\delta \lambda_{j_0^n}^{\tilde{j}_0^n} \epsilon_n^3 v_z(\tilde{s}^{\epsilon_n}, \tilde{z}^{\epsilon_n}) + o(\epsilon_n^3), \end{aligned}$$

which leads to a contradiction when n goes to $+\infty$.

Step 4: From (4.5) and Proposition 3.2, we deduce that in our domain of interest, the function $\Psi^{\epsilon, \delta}$ is actually smooth, and is therefore a legitimate test function for the second order operator of the dynamic programming equation. We then obtain from the supersolution property of v^ϵ that

$$\left(\beta v^\epsilon - \mathcal{L}\Psi^{\epsilon, \delta} - \tilde{U}(\Psi_x^{\epsilon, \delta}) \right) (\tilde{s}^\epsilon, \tilde{x}^\epsilon, \tilde{y}^\epsilon) \geq 0. \quad (4.8)$$

Moreover, by Step 3 and the continuity of $(s, z) \mapsto \mathcal{O}(s, z)$ in Assumption 3.4, the sequence $(\tilde{\xi}^\epsilon)_\epsilon$ is bounded. By classical results in the theory of viscosity solutions, there exists a sequence $\epsilon_n \rightarrow 0$ and some $\tilde{\xi}$ such that

$$(s_n, z_n, \xi_n) := (\tilde{s}^{\epsilon_n}, \tilde{z}^{\epsilon_n}, \tilde{\xi}^{\epsilon_n}) \longrightarrow (s_0, z_0, \tilde{\xi}).$$

Now recall that the function w is smooth in this case and that the function $\Psi^{\epsilon, \delta}$ has exactly the form given in Section 4.2. By the remainder estimate in (4.8), we obtain

$$\begin{aligned} & \frac{1}{2} v_{zz}(s_n, z_n) \eta(s_n, z_n) |\sigma^T(s_n) \xi_n|^2 + \frac{1}{2} (1 + \delta) \text{Tr} \left[\alpha \alpha^T(s_n, z_n) w_{\xi\xi}(s_n, z_n, \xi_n) \right] - \mathcal{A}\phi(s_n, z_n) \\ & - \mathcal{A}\Phi^{\epsilon_n}(s_n, z_n) + \mathcal{R}^\epsilon(s, z, \xi) \geq 0. \end{aligned}$$

We still have no guarantee that w is C^2 at $\tilde{\xi}$. Therefore, we carefully estimate the term involving $w_{\xi\xi}$. Indeed, the equation satisfied by w yields

$$\begin{aligned} & a(s_n, z_n) - \mathcal{A}\phi(s_n, z_n) + \delta \left(a(s_n, z_n) - \frac{1}{2} v_{zz}(s_n, z_n) \eta(s_n, z_n) |\sigma^T(s_n) \xi_n|^2 \right) \\ & - \mathcal{A}\Phi^{\epsilon_n}(s_n, z_n) + \mathcal{R}^\epsilon(s, z, \xi) \geq 0. \end{aligned} \quad (4.9)$$

Notice that the estimate on the remainder of Section 4.2 still hold true if terms involving $w_{\xi\xi}$ are replaced by means of the first corrector equation. Since the map $(s, z) \mapsto a(s, z)$ is continuous by Assumption 3.4, and all derivatives of Φ^ϵ vanish at the origin, we may send ϵ to 0 in (4.9) and obtain

$$a(s_0, z_0) - \mathcal{A}\phi(s_0, z_0) + \delta \left(a(s_0, z_0) - \frac{1}{2} \eta(s_0, z_0) |\sigma^T(s_0) \tilde{\xi}|^2 \right) \geq 0. \quad (4.10)$$

Since $\tilde{\xi}$ is bounded uniformly in δ , we let δ go to zero in (4.10) to obtain the desired result. \square

4.4. Viscosity Supersolution Property

In this section we will prove the following result.

Proposition 4.2. *Under Assumptions 3.1, 3.2 and 3.4, u_* is a viscosity supersolution of the second corrector equation.*

To prove this result, we will use two lemmas which are given in the Appendix (see Lemmas A.1 and A.2), to which we refer for the notations. Since they are pure calculus and can be proved by simple (but tedious) calculations, we will omit their proofs.

Let $k : \mathbb{R}^d \rightarrow \mathbb{R}$ be a positive, even (i.e. $k(-x) = k(x)$ for all $x \in \mathbb{R}^d$), C^∞ function with support in the closed unit ball of \mathbb{R}^d and unit total mass. For all $m > 0$, we define

$$k^m(x) := \frac{1}{m^d} k\left(\frac{x}{m}\right) \quad \text{and} \quad \tilde{w}^m(\cdot, \xi) := \int_{\mathbb{R}^d} k^m(\zeta) w(\cdot, \xi - \zeta) d\zeta - \int_{\mathbb{R}^d} k^m(\zeta) w(\cdot, -\zeta) d\zeta.$$

Proof [Proposition 4.2]. Let $(s_0, z_0, \phi) \in (0, +\infty)^{d+1} \times C^2((0, +\infty)^{d+1})$ be such that

$$(u_* - \phi)(s_0, z_0) < (u_* - \phi)(s, z), \quad \text{for all } (s, z) \in (0, +\infty)^d \times (0, +\infty) \setminus \{(s_0, z_0)\}. \quad (4.11)$$

By the definition of viscosity supersolutions, we need to show that

$$\mathcal{A}\phi(s_0, z_0) - a(s_0, z_0) \geq 0. \quad (4.12)$$

We argue by contradiction and assume that

$$\mathcal{A}\phi(s_0, z_0) - a(s_0, z_0) < 0. \quad (4.13)$$

Then by the continuity of ϕ and a , for some $r_0 > 0$, we will have

$$\mathcal{A}\phi(s, z) - a(s, z) \leq 0 \quad \text{on } B_{r_0}(s_0, z_0) \quad \text{for some } r_0 > 0. \quad (4.14)$$

Step 1: This first step is devoted to defining the test function we will consider in the sequel. First of all, we know from Lemma 4.2 that there exists a sequence (s^ϵ, z^ϵ) which realizes the lim for \hat{u}^ϵ , that is to say

$$(s^\epsilon, z^\epsilon) \xrightarrow{\epsilon \downarrow 0} (s_0, z_0) \quad \text{and} \quad \hat{u}^\epsilon(s^\epsilon, z^\epsilon, 0) \xrightarrow{\epsilon \downarrow 0} u_*(s_0, z_0).$$

It follows then easily that $l_*^\epsilon := \hat{u}^\epsilon(s^\epsilon, z^\epsilon, 0) - \phi(s^\epsilon, z^\epsilon) \xrightarrow{\epsilon \downarrow 0} 0$, and

$$(x^\epsilon, y^\epsilon) = (z^\epsilon - \mathbf{y}(s^\epsilon, z^\epsilon) \cdot \mathbf{1}_d, \mathbf{y}(s^\epsilon, z^\epsilon)) \xrightarrow{\epsilon \downarrow 0} (x_0, y_0) := (z_0 - \mathbf{y}(s_0, z_0) \cdot \mathbf{1}_d, \mathbf{y}(s_0, z_0)).$$

We then choose ϵ_0 , depending on z_0, s_0 and ϕ , such that for all $\epsilon \leq \epsilon_0$, we have

$$|z^\epsilon - z_0| + |s^\epsilon - s_0| \leq \frac{r_0}{4}, \quad |l_*^\epsilon| \leq 1. \quad (4.15)$$

By the continuity of ϕ , we may also introduce a constant $c_0 > 0$ such that

$$\sup_{(s,z) \in \bar{B}_{r_0/2}(s_0, z_0)} \{\phi(s, z) + C(s, z)\} + 3 \leq c_0 \left(\frac{r_0}{8}\right)^4, \quad (4.16)$$

where $C(s, z)$ is the continuous function appearing in the estimates A.7. Notice that since ϕ and C are continuous, the supremum on the compact set $\bar{B}_{r_0/2}(s_0, z_0)$ above is indeed finite. This justifies the existence of c_0 . We now define

$$\varphi^\epsilon(s, z) := \phi(s, z) - c_0 \left(|z - z^\epsilon|^4 + |s - s^\epsilon|^4\right).$$

Then, for all $\epsilon \leq \epsilon_0$ and for any $(s, z) \in \bar{B}_{r_0/2}(s_0, z_0)$ that

$$|z - z^\epsilon| + |s - s^\epsilon| \geq \frac{r_0}{4},$$

and therefore,

$$|z - z^\epsilon|^4 + |s - s^\epsilon|^4 \geq \frac{1}{16} (|z - z^\epsilon| + |s - s^\epsilon|)^4 \geq \left(\frac{r_0}{8}\right)^4.$$

Hence, we have using (4.16) and (4.15)

$$\begin{aligned} \varphi^\epsilon(s, z) + I_\epsilon^* + C(s, z) &= \phi(s, z) + C(s, z) - c_0 \left(|z - z^\epsilon|^4 + |s - s^\epsilon|^4\right) + I_\epsilon^* \\ &\leq c_0 \left(\frac{r_0}{8}\right)^4 - 3 - c_0 \left(|z - z^\epsilon|^4 + |s - s^\epsilon|^4\right) + 1 \\ &\leq c_0 \left(\left(\frac{r_0}{8}\right)^4 - |z - z^\epsilon|^4 - |s - s^\epsilon|^4\right) - 2 \leq -2, \end{aligned} \quad (4.17)$$

whenever $(s, z) \in \partial B_{r_0/2}(s_0, z_0)$.

Before defining our final test function, we provide another parameter. Throughout the rest of the proof, we let $m \in (0, 1]$. Now for any $\delta \in (0, 1)$ and $v > 0$, let $h^{\delta, v}$ be the function defined by Lemma A.2, and introduce a parameter $\xi^* := 1 \vee \tilde{\xi}_0 \vee \tilde{\xi}^*$, where $\tilde{\xi}_0 > 0$ is greater than $\eta^s(s_0, z_0)$ times the diameter of $\mathcal{O}(s_0, z_0)$ and large enough so that for every $|\xi| \geq \tilde{\xi}_0$, $\tilde{w}_{\xi\xi}^m(s, z, \xi) = 0$, for every $(s, z) \in B_{r_0/2}(s_0, z_0)$, and $\tilde{\xi}^*$ is such that for any $\xi \in B_{\tilde{\xi}^*}(0)^c$ and $(s, z) \in \bar{B}_{r_0/2}(s_0, z_0)$, we have

$$-\frac{1}{2} v_{zz} |\sigma^T \xi|^2 - \mathcal{A}(\varphi^\epsilon + \phi) > \frac{1}{2} \text{Tr}[\alpha \alpha^T] C(s, z) \left(C^* + \frac{\sqrt{d} \delta \underline{\lambda}}{3L}\right) + 1, \quad (4.18)$$

where C^* is the constant introduced in Lemma A.2 and $C(s, z)$ is the function introduced in Lemma A.1 (we remind the reader that they are both uniform in m).

Finally, define the C^2 test function $\psi^{\epsilon, \delta, m}(s, x, y)$ and the corresponding $\hat{\psi}^{\epsilon, \delta, m}(s, z, \zeta)$ by

$$\begin{aligned}\hat{\psi}^{\epsilon, \delta, m}(s, z, \zeta) &:= v(s, z) - \epsilon^2 \varphi^\epsilon(s, z) - \epsilon^2 l_\epsilon^* - \epsilon^4 (1 - \delta) \tilde{w}^m(s, z, \zeta) H(\zeta), \\ H(\zeta) &:= h^{\delta, v} \left(\frac{\zeta}{\zeta^*} \right).\end{aligned}$$

Step 2: In this step, we modify the test function once again in order to recover the interior maximizer property. We first compute that:

$$\begin{aligned}I^{\epsilon, \delta, m}(s, z, \zeta) &:= \epsilon^{-2} (\hat{v}^\epsilon - \hat{\psi}^{\epsilon, \delta, m})(s, z, \zeta) \\ &= \varphi^\epsilon(s, z) - \hat{u}^\epsilon(s, z, \zeta) + l_\epsilon^* - \epsilon^2 w(s, z, \zeta) + \epsilon^2 (1 - \delta) H(\zeta) \tilde{w}^m(s, z, \zeta).\end{aligned}\tag{4.19}$$

In particular, this implies

$$I^{\epsilon, \delta, m}(s^\epsilon, z^\epsilon, 0) = \epsilon^2 (1 - \delta) \tilde{w}^m(s^\epsilon, z^\epsilon, 0),$$

which, according to Lemma A.1, goes to 0 as ϵ goes to 0, uniformly in $m \in (0, 1]$. Therefore, we have for ϵ small enough

$$I^{\epsilon, \delta, m}(s^\epsilon, z^\epsilon, 0) \geq -1.\tag{4.20}$$

Furthermore, since $v^\epsilon \leq v$, we also have easily

$$I^{\epsilon, \delta, m}(s, z, \zeta) \leq \varphi^\epsilon(s, z) + l_\epsilon^* + \epsilon^2 (1 - \delta) H(\zeta) \tilde{w}^m(s, z, \zeta).\tag{4.21}$$

Now, we use the fact that

$$0 \leq \delta \leq 1, v_z(s, z) > 0, 0 \leq H(\zeta) \leq \mathbf{1}_{|\zeta| \leq a^\delta \zeta^*} \quad \text{and} \quad 0 \leq \tilde{w}^m(s, z, \zeta) \leq 2C(s, z)(1 + |\zeta|),$$

in (4.21) to obtain

$$I^{\epsilon, \delta, m}(s, z, \zeta) \leq \varphi^\epsilon(s, z) + l_\epsilon^* + 2\epsilon^C(s, z)(1 + a^\delta \zeta^*) \leq \varphi^\epsilon(s, z) + l_\epsilon^* + C(s, z),$$

provided that $\epsilon \leq \epsilon^\delta := (2(1 + a^\delta \zeta^*))^{1/2}$.

Define the set $Q_{s_0, z_0} := \{(s, z, \zeta), (s, z) \in \bar{B}_{r_0}(s_0, z_0)\}$. Let us then distinguish two cases. First, we assume that $(s, z, \zeta) \in \partial Q_{s_0, z_0}$ for every ζ . Then, if we take $\epsilon \leq \epsilon_0 \wedge \epsilon^\delta$, using (4.17) and (4.22), we obtain that for any ζ

$$I^{\epsilon, \delta, m}(z, \zeta) \leq -2.\tag{4.22}$$

We assume next that $(s, z, \zeta) \in \text{int}(Q_{s_0, z_0})$. Then, once again for $\epsilon \leq \epsilon^\delta \wedge \epsilon_0$, we have

$$I^{\epsilon, \delta, m}(z, \zeta) \leq \varphi^\epsilon(s, z) + l_\epsilon^* + v_z(s, z) \leq C(s_0, z_0) < +\infty,\tag{4.23}$$

for some constant $C(s_0, z_0)$ depending only on ϕ , s_0 and z_0 , since (s, z) lies in a compact set, $l_\epsilon^* \leq 1$ and all the functions appearing on the right-hand side of (4.23) are continuous. This implies that

$$v(\epsilon, \delta, m) := \sup_{(s, z, \zeta) \in Q_{s_0, z_0}} I^{\epsilon, \delta, m}(s, z, \zeta) < +\infty.$$

By definition, we can therefore for each $n \geq 1$ find $(\hat{s}_n, \hat{z}_n, \hat{\zeta}_n) \in \text{int}(Q_{s_0, z_0})$ such that

$$I^{\epsilon, \delta, m}(\hat{s}_n, \hat{z}_n, \hat{\zeta}_n) \geq v(\epsilon, \delta, m) - \frac{1}{2n}. \quad (4.24)$$

Since we have no guarantee that the maximizer above exists, we modify once more our test function. Let f be an even smooth function such that $f(0) = 1$, $f(x) = 0$ if $|x| \geq 1$ and $0 \leq f \leq 1$. We then define the test function $\Psi^{\epsilon, \delta, m, n}(s, x, y)$ and the corresponding

$$\hat{\Psi}^{\epsilon, \delta, m, n}(s, z, \zeta) := \hat{\psi}^{\epsilon, \delta, m}(s, z, \zeta) - \frac{\epsilon^2}{n} f(|\zeta - \hat{\zeta}_n|).$$

Consider then

$$I^{\epsilon, \delta, m, n}(s, z, \zeta) := \epsilon^{-2} \left(\hat{\psi}^\epsilon - \hat{\psi}^{\epsilon, \delta, m, n} \right) (s, z, \zeta) = I^{\epsilon, \delta, m}(s, z, \zeta) + \frac{1}{n} f(|\zeta - \hat{\zeta}_n|).$$

Notice now that for any $(s, z, \zeta) \in Q_{s_0, z_0}$, we have using (4.24)

$$I^{\epsilon, \delta, m, n}(\hat{s}_n, \hat{z}_n, \hat{\zeta}_n) = I^{\epsilon, \delta, m}(\hat{s}_n, \hat{z}_n, \hat{\zeta}_n) + \frac{1}{n} \geq v(\epsilon, \delta, m) + \frac{1}{2n} \geq I^{\epsilon, \delta, m}(s, z, \zeta) + \frac{1}{2n}. \quad (4.25)$$

Moreover, by the definition of f , we have:

$$I^{\epsilon, \delta, m, n}(s, z, \zeta) = I^{\epsilon, \delta, m}(s, z, \zeta) \quad \text{if } \zeta \in B_1(\hat{\zeta}_n)^c \quad \text{and } (s, z, \zeta) \in Q_{s_0, z_0}.$$

This equality and (4.25) imply

$$\sup_{(s, z, \zeta) \in Q_{s_0, z_0}} I^{\epsilon, \delta, m, n}(s, z, \zeta) = \sup_{(s, z, \zeta) \in \bar{B}_1(\hat{\zeta}_n) \cap Q_{s_0, z_0}} I^{\epsilon, \delta, m, n}(s, z, \zeta).$$

Since $\bar{B}_1(\hat{\zeta}_n) \cap Q_{s_0, z_0}$ is a compact set, we deduce that there exists some $(s_n, z_n, \zeta_n) \in Q_{s_0, z_0}$ which maximizes $I^{\epsilon, \delta, m, n}$. We now claim that we actually have $(s_n, z_n, \zeta_n) \in \text{int}(Q_{s_0, z_0})$. Indeed, we have by (4.20)

$$I^{\epsilon, \delta, m, n}(s_n, z_n, \zeta_n) \geq I^{\epsilon, \delta, m, n}(s^\epsilon, z^\epsilon, 0) \geq I^{\epsilon, \delta, m}(s^\epsilon, z^\epsilon, 0) = 0,$$

and, by (4.22),

$$I^{\epsilon, \delta, m, n}(s, z, \zeta) \leq I^{\epsilon, \delta, m}(s, z, \zeta) + \frac{1}{n} \leq -2 + \frac{1}{n} < 0, \quad (s, z, \zeta) \in \partial Q_{s_0, z_0}$$

for $(s, z, \zeta) \in \partial Q_{s_0, z_0}$.

By the viscosity subsolution property of v^ϵ at the point (s_n, z_n, ξ_n) , with corresponding (s_n, x_n, y_n) , it follows that

$$\min_{(i,j) \in \mathcal{F}} \left\{ \beta v^\epsilon - \mathcal{L} \Psi^{\epsilon, \delta, m, n} - \tilde{U}(\Psi_x^{\epsilon, \delta, m, n}), \quad \Lambda_{i,j}^\epsilon \cdot (\Psi_x^{\epsilon, \delta, m, n}, \Psi_y^{\epsilon, \delta, m, n}) \right\} \leq 0. \quad (4.26)$$

Step 3: Our aim in this step is to show that for ϵ small enough and n large enough, we have:

$$D^{i,j} := \Lambda_{i,j}^\epsilon \cdot (\Psi_x^{\epsilon, \delta, m, n}, \Psi_y^{\epsilon, \delta, m, n})(s_n, x_n, y_n) > 0 \quad \text{for all } (i, j) \in \mathcal{F}. \quad (4.27)$$

We easily compute for $0 \leq i \leq d$, with the convention that $y_0 = x$ and $e_0 = 0$, that:

$$\begin{aligned} \Psi_{y_i}^{\epsilon, \delta, m, n}(s, x, y) &= v_z(s, z) - \epsilon^2 \varphi_z^\epsilon(s, z) - \epsilon^3 (1 - \delta) (w^m H)_\xi(s, z, \xi) \cdot (e_i - \mathbf{y}_z(s, z)) \\ &\quad - \epsilon^4 (1 - \delta) (w_z^m H)(s, z, \xi) - \frac{\epsilon f'(|\xi - \hat{\xi}_n|)}{n |\xi - \hat{\xi}_n|} \xi \cdot (e_i - \mathbf{y}_z(s, z)). \end{aligned}$$

and

$$D^{i,j} = \epsilon^3 \left[\lambda^{i,j} v_z(s_n, z_n) - (1 - \delta) (w^m H)_\xi(s_n, z_n, \xi_n) \cdot (e_i - e_j) \right] - E^\epsilon - F^{\epsilon, n},$$

where

$$\begin{aligned} E^\epsilon &:= \lambda^{i,j} \left[\epsilon^5 (\phi_z(s_n, z_n) - 4C(z_n - z^\epsilon)^3) + \epsilon^6 (1 - \delta) (w^m H)_\xi(s_n, z_n, \xi_n) \cdot (e_i - \mathbf{y}_z(s_n, z_n)) \right] \\ &\quad + \lambda^{i,j} \epsilon^7 (1 - \delta) (w_z^m H)(s_n, z_n, \xi_n), \\ F^{\epsilon, n} &:= \frac{\epsilon f'(|\xi_n - \hat{\xi}_n|)}{n |\xi_n - \hat{\xi}_n|} \xi_n \cdot (e_i - e_j + \lambda^{i,j} \epsilon^3 (e_i - \mathbf{y}_z(s_n, z_n))). \end{aligned}$$

Recall that $\xi^* \geq 1$. Then, from Lemma A.2, we have

$$0 \leq H \leq 1, \quad |H_\xi| \leq \frac{\sqrt{d} v \delta}{3 \xi^*} \leq \frac{\sqrt{d} v \delta}{3} \quad \text{and} \quad H = 0 \quad \text{for} \quad |\xi| \geq a^\delta \xi^*,$$

we deduce

$$\begin{aligned} |E^\epsilon| &\leq \lambda^{i,j} \epsilon^5 \left[\phi_z(s_n, z_n) + 4c_0 |z_n - z^\epsilon|^3 + \epsilon C(s_0, z_0) (|w_\xi^m| H + w^m |H_\xi|) (s_n, z_n, \xi_n) \right. \\ &\quad \left. + \epsilon^2 L v_z(s_n, z_n) |\xi_n| \mathbf{1}_{H(\xi_n) > 0} \right] \\ &\leq C(s_0, z_0) \epsilon^5 [1 + \epsilon (1 + a^\delta) + \epsilon^2 a^\delta \xi^*], \end{aligned} \quad (4.28)$$

for some constant $C(s_0, z_0)$ which can change value from line to line. Then we also have easily for some constant denoted Const , which can also change value from line to line

$$|F^{\epsilon, n}| \leq \text{Const} \frac{\epsilon}{n}. \quad (4.29)$$

Let us now study the term

$$\begin{aligned} G^\epsilon &:= \lambda^{i,j} v_z - (1 - \delta)(w^m H)_{\xi_i} (e_i - e_j) \\ &= \lambda^{i,j} v_z - (1 - \delta)(w_{\xi_i}^m - w_{\xi_j}^m) H - (1 - \delta) w^m (H_{\xi_i} - H_{\xi_j}), \end{aligned}$$

where we suppressed the dependence in (s_n, z_n, ξ_n) for simplicity. Using Lemma A.1(iii) and the fact that w^m and H are positive and $w^m \leq L v_z |\xi|$

$$\begin{aligned} G^\epsilon &\geq \lambda^{i,j} v_z - \lambda^{i,j} (1 - \delta) v_z - (1 - \delta) L v_z |\xi_n| \left(|H_{\xi_i}| + |H_{\xi_j}| \right) \\ &\geq \lambda^{i,j} v_z \left(\delta - \frac{L |\xi_n|}{\lambda^{i,j}} (1 - \delta) \left(|H_{\xi_i}| + |H_{\xi_j}| \right) \right) \\ &\geq \lambda^{i,j} v_z \left(\delta - \frac{L |\xi_n|}{\underline{\lambda}} (1 - \delta) \left(|H_{\xi_i}| + |H_{\xi_j}| \right) \right). \end{aligned}$$

But by (iii) of Lemma A.2, we know that for all $0 \leq i, j \leq d$, $|\xi| |H_{\xi_i}(\xi)| \leq v \delta$. Fix $v := \underline{\lambda}/(2L)$, we deduce

$$G^\epsilon \geq \lambda^{i,j} \delta^2 v_z. \quad (4.30)$$

Finally, we have obtained

$$D^{i,j} \geq \lambda^{i,j} \delta^2 v_z \epsilon^3 - C(s_0, z_0) \epsilon^5 \left(1 + \epsilon (1 + a^\delta) + \epsilon^2 a^\delta \xi^* \right) - \text{Const} \frac{\epsilon}{n}.$$

Next, there is by hypothesis some constant $\tilde{C} > 0$ such that $v_z \geq \tilde{C}$, and therefore there is a $\tilde{\epsilon}^\delta$ such that for all $\epsilon \leq \tilde{\epsilon}^\delta$:

$$C(s_0, z_0) \epsilon^5 \left(1 + \epsilon (1 + a^\delta) + \epsilon^2 a^\delta \xi^* \right) \leq \frac{\lambda^{i,j} \delta^2 \tilde{C} \epsilon^3}{4}.$$

Then, for all $n \geq N_{\epsilon, \delta} := \frac{4 \text{Const}}{\epsilon^2 \tilde{C} \delta^2}$, we have

$$\text{Const} \frac{\epsilon}{n} \leq \frac{\lambda^{i,j} \delta^2 \tilde{C} \epsilon^3}{4}.$$

We then conclude that, for $\epsilon \leq \tilde{\epsilon}^\delta$ and $n \geq N_{\epsilon, \delta}$, we have $D^{i,j} \geq \lambda^{i,j} \delta^2 \tilde{C} \epsilon^3 / 2 > 0$, and by the arbitrariness of $i, j = 0, \dots, d$, we deduce from (4.26) that

$$J^{\epsilon, \delta, m, n} := \frac{1}{\epsilon^2} \left[\beta v^\epsilon - \mathcal{L} \Psi^{\epsilon, \delta, m, n} - \tilde{U} \left(\Psi_x^{\epsilon, \delta, m, n} \right) \right] (s_n, x_n, y_n) \leq 0. \quad (4.31)$$

Step 4: We now consider the remainder estimate. Using the general expansion result, we have

$$J^{\epsilon, \delta, m, n} = (-v_{zz}) \frac{|\sigma^T \xi_n|^2}{2} + \frac{1 - \delta}{2} \text{Tr} \left[\alpha \alpha^T (\tilde{w}^m H)_{\xi \xi} \right] - \mathcal{A}(\phi + \varphi^\epsilon) + \mathcal{R}, \quad (4.32)$$

where

$$\mathcal{R} = \mathcal{R}_{\tilde{U}} + \mathcal{R}_\phi + \mathcal{R}_f + \mathcal{R}_{wH},$$

and using the same calculations as in the remainder estimate of Section 4.2

$$|\mathcal{R}_{\tilde{U}}| = \epsilon^{-2} \left| \tilde{U}(\Psi_x^{\epsilon, \delta, m, n}) - \tilde{U}(v_z) - \epsilon^2 \tilde{U}'(\phi_z) \right| \leq \text{Const} \left(\epsilon + |\epsilon \xi_n| + \frac{1}{n} \right)$$

$$|\mathcal{R}_\phi| \leq \text{Const} \left[|\epsilon \xi_n| + |\epsilon \xi_n|^2 \right]$$

$$|\mathcal{R}_f| \leq \frac{\text{Const}}{n}$$

$$|\mathcal{R}_{wH}| \leq \text{Const} \left(1 + |\epsilon \xi_n|^2 \right) \left(\epsilon + \epsilon^2 |\xi_n| + \epsilon |\xi_n| \right) \mathbf{1}_{|\xi_n| \leq a^\delta \xi^*}.$$

We now prove that (ξ_n) is bounded uniformly in ϵ , so that $\epsilon \xi_n \rightarrow 0$ and $\epsilon \rightarrow 0$. First, we estimate that:

$$\begin{aligned} |(\tilde{w}^m H)_{\xi\xi}|(s, z, \xi) &\leq (|\tilde{w}_{\xi\xi}^m| |H| + 2|\tilde{w}_\xi^m| |H_\xi| + |\tilde{w}^m| |H_{\xi\xi}|)(s, z, \xi) \\ &\leq C(s, z) \left(1 + 2\frac{\sqrt{d}v\delta}{3} + (1 + |\xi|) |H_{\xi\xi}| \right) \\ &\leq C(s, z) \left(1 + 2\frac{\sqrt{d}v\delta}{3} + C^* \right) \leq \tilde{C}(s, z), \end{aligned}$$

where we used A.2 and where $\tilde{C}(s, z)$ depends only on (s, z) (and not on δ). From (4.31), we deduce that

$$\begin{aligned} (-v_{zz}(s_n, z_n)) \frac{|\sigma^T \xi_n|^2}{2} + \mathcal{R} &\leq \left(\mathcal{A}\phi + |\alpha\alpha^T| \tilde{C} \right) (s_n, z_n) \leq \sup_{(s, z) \cap Q_{s_0, z_0}} \left(\mathcal{A}\phi + |\alpha\alpha^T| \tilde{C} \right) \\ &\leq \text{Const}. \end{aligned}$$

Using the above estimates for \mathcal{R} , we deduce that

$$(-v_{zz}) \frac{|\sigma^T \xi_n|^2}{2} - \text{Const} \left(\epsilon + |\epsilon \xi_n| (1 + \epsilon) + |\epsilon \xi_n|^2 (1 + \epsilon (1 + a^\delta (1 + \epsilon))) + \frac{1}{n} \right) \leq \text{Const},$$

from which we deduce that for some constant C_0 depending on (s_0, z_0)

$$C_0 |\xi_n|^2 - \text{Const} \left(\epsilon + |\epsilon \xi_n| (1 + \epsilon) + |\epsilon \xi_n|^2 (1 + \epsilon (1 + a^\delta (1 + \epsilon))) + \frac{1}{n} \right) \leq \text{Const}. \quad (4.33)$$

Let us now assume that ξ_n is unbounded as ϵ goes to 0, that is to say that we can find some sequence ϵ_k going to 0, such that $|\xi_n|$ goes to $+\infty$ as k goes to $+\infty$. Then, taking k and n large enough contradicts (4.33). Therefore ξ_n is bounded by some $\hat{\xi}^\delta$ depending only on δ .

We deduce from all these estimates that there is a $\hat{\epsilon}^\delta$ and a \tilde{N}_ϵ such that if $\epsilon \leq \hat{\epsilon}^\delta$ and $n \geq \tilde{N}_\epsilon$, we have

$$|\mathcal{R}| \leq 1.$$

Step 5: For n greater than all the previously introduced N 's and ϵ smaller than all the ones previously introduced, we now show that $|\xi_n| \leq \xi^*$.

We argue by contradiction and we suppose that $|\xi_n| > \xi^*$. We remind the reader that $\xi^* \geq 1$ and that $|w_\xi^m| \leq L$. Then we know that $w_{\xi\xi}^m(\cdot, \xi_n) = 0$. By (4.31), the expansion (4.32), the result of Lemma A.2(iii),(iv), we see that:

$$\begin{aligned} (-v_{zz}(s_n, z_n)) \frac{|\sigma^T \xi_n|^2}{2} - \mathcal{A}(\phi + \varphi^\epsilon) &\leq -\frac{1-\delta}{2} \text{Tr} [\alpha \alpha^T (s_n, z_n) (\tilde{w}^m H)_{\xi\xi}(s_n, z_n, \xi_n)] - \mathcal{R} \\ &\leq -\frac{1-\delta}{2} \text{Tr} [\alpha \alpha^T (s_n, z_n) (\tilde{w}^m H_{\xi\xi} + 2\tilde{w}_\xi^m H_\xi)] + 1 \\ &\leq C(s_n, z_n) \text{Tr} [\alpha \alpha^T (s_n, z_n)] \left(C^* + \frac{\sqrt{d}\delta}{3\underline{\lambda}} \right) + 1, \end{aligned}$$

contradicting (4.18). Hence $(\xi_n)_n$ is bounded by ξ^* (which does not depend on ϵ , δ , m or n). In particular, this implies that the function H applied to ξ_n is always equal to 1. Therefore, by the boundedness of (s_n, z_n, ξ_n) and by classical results on the theory of viscosity solutions, there exists some $\bar{\xi}$ such that by letting n go to $+\infty$ and then ϵ to 0 (along some subsequence if necessary) in (4.32), we obtain by using Lemma A.1 (iv):

$$\begin{aligned} 0 &\geq -v_{zz}(s_0, z_0) \frac{|\sigma^T(s_0) \bar{\xi}|^2}{2} + \frac{(1-\delta)}{2} \text{Tr} [\alpha \alpha^T (s_0, z_0) \tilde{w}_{\xi\xi}^m(s_0, z_0, \bar{\xi})] - \mathcal{A}\phi(s_0, z_0) \\ &\geq (1-\delta)a(s_0, z_0) - \delta v_{zz}(s_0, z_0) \frac{|\sigma^T(s_0) \bar{\xi}|^2}{2} - \mathcal{A}\phi(s_0, z_0) \\ &\quad + \frac{1-\delta}{2} v_{zz}(s_0, z_0) \int_{\mathbb{R}^d} k^m(\zeta) (|\sigma^T(s_0)(\bar{\xi} - \zeta)|^2 - |\sigma^T(s_0) \bar{\xi}|^2) d\zeta. \end{aligned} \quad (4.34)$$

Now using the fact that the function k^m is even, we have

$$\int_{\mathbb{R}^d} k^m(\zeta) (|\sigma^T(s_0)(\bar{\xi} - \zeta)|^2 - |\sigma^T(s_0) \bar{\xi}|^2) d\zeta = \int_{\mathbb{R}^d} k^m(\zeta) |\sigma^T(s_0) \zeta|^2 d\zeta.$$

Since $\int_{\mathbb{R}^d} k^m(\zeta) |\sigma^T(s_0) \zeta|^2 d\zeta \xrightarrow{m \rightarrow 0} 0$, and $\bar{\xi}$ is uniformly bounded in m and δ , we can let δ and m go to 0 in (4.34) to obtain

$$\mathcal{A}\phi(s_0, z_0) - a(s_0, z_0) \geq 0,$$

which is the required contradiction to (4.13), and completes the proof of the required result (4.12). \square

5. Wellposedness of the First Corrector Equation

In this section, we collect the main proofs which allow us to obtain the wellposedness of the first corrector equation (2.6). Since the variables (s, z) are

frozen in this equation, we simplify the notations by suppressing the dependence on them. Recall that since the set C is bounded, convex and closed, the supremum in the definition of the convex function δ_C is always attained at the boundary ∂C . Moreover $0 \in \text{int}(C)$, we may find two constants $L, L' > 0$ such that

$$L'|\rho| \leq \delta_C(\rho) \leq L|\rho|. \quad (5.1)$$

5.1. Uniqueness and Comparison

Proof of Theorem 3.1. Fix some $(\epsilon, \nu) \in (0, 1) \times (0, +\infty)$ and define for $(\rho, y) \in \mathbb{R}^d \times \mathbb{R}^d$

$$\bar{w}^\epsilon(\rho, y) := (1 - \epsilon)w_1(\rho) - w_2(y), \quad \phi_\nu(\rho, y) = \frac{1}{2\nu}|\rho - y|^2.$$

Since w_1 is a viscosity subsolution of (2.6), then its gradient takes values in C in the viscosity sense, which implies that w_1 is L -Lipschitz. Then, for $y \neq 0$:

$$\begin{aligned} \bar{w}^\epsilon(\rho, y) - \phi_\nu(\rho, y) &= (1 - \epsilon)(w_1(\rho) - w_1(y)) - \frac{1}{2\nu}|\rho - y|^2 + (1 - \epsilon)w_1(y) - w_2(y) \\ &\leq (1 - \epsilon)L|\rho - y| - \frac{1}{2\nu}|\rho - y|^2 + (1 - \epsilon)w_1(y) - w_2(y) \\ &\leq \frac{(1 - \epsilon)^2 L^2 \nu}{2} + (1 - \epsilon)w_1(y) - w_2(y) \\ &= \frac{(1 - \epsilon)^2 L^2 \nu}{2} + \delta_C(y) \left((1 - \epsilon) \frac{w_1(y)}{\delta_C(y)} - \frac{w_2(y)}{\delta_C(y)} \right). \end{aligned}$$

By the growth conditions on w_1, w_2 , together with (5.1), this implies that:

$$\lim_{|(\rho, y)| \rightarrow +\infty} \bar{w}^\epsilon(\rho, y) - \phi_\nu(\rho, y) = -\infty.$$

Then, the difference $\bar{w}^\epsilon - \phi_\nu$ has a global maximizer $(\rho^{\epsilon, \nu}, y^{\epsilon, \nu}) \in \mathbb{R}^d \times \mathbb{R}^d$ satisfying the lower bound

$$(\bar{w}^\epsilon - \phi_\nu)(\rho^{\epsilon, \nu}, y^{\epsilon, \nu}) \geq (\bar{w}^\epsilon - \phi_\nu)(0, 0) = 0. \quad (5.2)$$

By the Crandall-Ishii Lemma (see Theorem 3.2 in [10]), it follows that for any $\eta > 0$, there exist symmetric positive matrices X and Y such that

$$\begin{aligned} (D_\rho \phi_\nu(\rho^{\epsilon, \nu}, y^{\epsilon, \nu}), X) &= \left(\frac{\rho^{\epsilon, \nu} - y^{\epsilon, \nu}}{\nu}, X \right) \in \bar{J}^{2,+}((1 - \epsilon)w_1)(\rho^{\epsilon, \nu}) \\ (-D_y \phi_\nu(\rho^{\epsilon, \nu}, y^{\epsilon, \nu}), Y) &= \left(\frac{\rho^{\epsilon, \nu} - y^{\epsilon, \nu}}{\nu}, Y \right) \in \bar{J}^{2,-}w_2(y^{\epsilon, \nu}), \end{aligned} \quad (5.3)$$

and

$$\begin{pmatrix} X & 0 \\ 0 & -Y \end{pmatrix} \leq A + \eta A^2, \quad \text{with } A := D^2 \phi_\nu(\rho^{\epsilon, \nu}, y^{\epsilon, \nu}) = \frac{1}{\nu} \begin{pmatrix} I_d & -I_d \\ -I_d & I_d \end{pmatrix}.$$

The above matrix inequality directly implies that $X \leq Y$. We now use (5.3) to arrive at

$$\frac{\rho^{\epsilon,v} - y^{\epsilon,v}}{(1-\epsilon)v} \in \bar{J}^{1,+} w_1(\rho^{\epsilon,v}).$$

In addition, since w_1 is a viscosity subsolution, $Dw_1 \in C$ in the viscosity sense. This implies that for all $0 \leq i, j \leq d$,

$$-\lambda^{i,i} \leq \frac{\rho_i^{\epsilon,v} - y_i^{\epsilon,v}}{(1-\epsilon)v} - \frac{\rho_j^{\epsilon,v} - y_j^{\epsilon,v}}{(1-\epsilon)v} \leq \lambda^{i,j}.$$

Since $\epsilon \in (0, 1)$, we have

$$-\lambda^{i,i} < \frac{\rho_i^{\epsilon,v} - y_i^{\epsilon,v}}{v} - \frac{\rho_j^{\epsilon,v} - y_j^{\epsilon,v}}{v} < \lambda^{i,j}, \quad \text{and therefore } \frac{\rho^{\epsilon,v} - y^{\epsilon,v}}{v} \in \text{int}(C).$$

Consequently, it follows from (5.3) and the viscosity subsolution and supersolution of w_1 and w_2 that

$$-\frac{|\sigma\rho^{\epsilon,v}|^2}{2} - \frac{1}{2(1-\epsilon)} \text{Tr} [\bar{\alpha}\bar{\alpha}^T X] + a_1 \leq 0 \leq -\frac{|\sigma y^{\epsilon,v}|^2}{2} - \frac{1}{2} \text{Tr} [\bar{\alpha}\bar{\alpha}^T Y] + a_2.$$

Since $X \leq Y$, this provides:

$$(1-\epsilon)a_1 - a_2 \leq \frac{1}{2} \text{Tr} [\bar{\alpha}\bar{\alpha}^T (X - Y)] + (1-\epsilon) \frac{|\sigma\rho^{\epsilon,v}|^2}{2} - \frac{|\sigma y^{\epsilon,v}|^2}{2} \leq \frac{|\sigma\rho^{\epsilon,v}|^2}{2} - \frac{|\sigma y^{\epsilon,v}|^2}{2}. \quad (5.4)$$

We now show that $(y^{\epsilon,v}, \rho^{\epsilon,v})_v$ remains bounded as v tends to zero. We argue by contradiction, assuming to the contrary that

$$|y^{\epsilon,v_n}| \longrightarrow \infty \quad \text{for some sequence } v_n \longrightarrow \infty.$$

Since w_1 is Lipschitz, this implies that

$$(\bar{w}^\epsilon - \phi_{v_n})(\rho^{\epsilon,v_n}, y^{\epsilon,v_n}) \leq \frac{((1-\epsilon)L)^2 v_n}{2} + \delta_C(y^{\epsilon,v_n}) \frac{1-\epsilon}{\delta_C(y^{\epsilon,v_n})} (w_1(y^{\epsilon,v_n}) - w_2(y^{\epsilon,v_n})).$$

Arguing as in the beginning of this proof, we see that $(\bar{w}^\epsilon - \phi_{v_n})(\rho^{\epsilon,v_n}, y^{\epsilon,v_n}) \longrightarrow -\infty$, contradicting (5.2).

Similarly, using the normalization $w_1(0) = 0$, we have

$$(\bar{w}^\epsilon - \phi_{v_n})(\rho^{\epsilon,v_n}, y^{\epsilon,v_n}) \leq (1-\epsilon)L|\rho^{\epsilon,v_n}| - w_2(y^{\epsilon,v_n}) - \frac{|\rho^{\epsilon,v_n} - y^{\epsilon,v_n}|^2}{v_n}.$$

Since $(y^{\epsilon,v_n})_n$ was just shown to be bounded, we see that $|\rho^{\epsilon,v_n}| \longrightarrow \infty$ implies $(\bar{w}^\epsilon - \phi_{v_n})(\rho^{\epsilon,v_n}, y^{\epsilon,v_n}) \longrightarrow -\infty$, and thus a contradiction.

By standard techniques from the theory of viscosity solutions, we may then construct a $\rho^\epsilon \in \mathbb{R}^d$ and a sequence $(\tilde{v}_n)_{n \geq 0}$ converging to zero such that

$(\rho^{\epsilon, \bar{v}_n}, y^{\epsilon, \bar{v}_n}) \rightarrow (\rho^\epsilon, \rho^\epsilon)$, as $n \rightarrow \infty$. Passing to the limit in (5.4) along this sequence, we see that $(1 - \epsilon)a_1 - a_2 \leq 0$, which implies that $a_1 \leq a_2$ by the arbitrariness of $\epsilon \in (0, 1)$. \square

The following uniqueness result is an immediate consequence.

Corollary 5.1. *There is at most one $a \in \mathbb{R}$ such that (2.6) has a viscosity solution \bar{w} satisfying the growth condition $\bar{w}(\rho)/\delta_C(\rho) \rightarrow 1$, as $|\rho| \rightarrow \infty$.*

Remark 5.1.

- (i) In the context of [21], C is just the closed unit ball, then it is clear that $\delta_C(\rho) = |\rho|$, and the growth condition in the previous result reduces to $\bar{w}(\rho)/|\rho| \rightarrow 1$, as $|\rho| \rightarrow \infty$.
- (ii) In the one-dimensional case $d = 1$, we directly compute that $\delta_C(\rho) = \lambda^{1,0}\rho^+ + \lambda^{0,1}\rho^-$. So the above growth condition is the sharpest one for the explicit solution of (2.6) given in Section 4 of [40]. \square

5.2. Optimal Control Approximation of the First Corrector Equation

In ergodic control, it is standard to introduce an approximation by a sequence of infinite horizon standard control problems with a vanishing discount factor $\eta > 0$:

$$\max_{(i,j) \in \mathcal{J}} \left\{ -\frac{|\sigma^T \rho|^2}{2} - \frac{1}{2} \text{Tr} [\bar{\alpha} \bar{\alpha}^T D^2 \bar{w}^\eta(\rho)] + \eta \bar{w}^\eta(\rho), -\lambda^{i,j} + \frac{\partial \bar{w}^\eta}{\partial \rho_i}(\rho) - \frac{\partial \bar{w}^\eta}{\partial \rho_j}(\rho) \right\} = 0, \quad (5.5)$$

together with the growth condition

$$\lim_{|\rho| \rightarrow +\infty} \frac{\bar{w}^\eta(\rho)}{\delta_C(\rho)} = 1. \quad (5.6)$$

The next result states the existence of a unique solution of the approximating control problem.

Proposition 5.1. *For every $\eta \in (0, 1]$, there is a unique viscosity solution \bar{w}^η of (5.5)–(5.6). Moreover, \bar{w}^η is L -Lipschitz (with a constant L independent of η), and we have the following estimate*

$$(\delta_C(\rho) - K_1)^+ \leq \bar{w}^\eta(\rho) \leq \frac{K_2}{\eta} + \delta_C(\rho), \quad \rho \in \mathbb{R}^d. \quad (5.7)$$

Proof. Uniqueness is a classical consequence of Theorem 3.1. We establish existence of a viscosity solution by an application of Perron's method, which requires to find appropriate sub and supersolutions. The remaining properties are immediate consequences. We then introduce:

$$\underline{w}(\rho) := (\delta_C(\rho) - K_1)^+, \quad \bar{\bar{w}}(\rho) := \frac{K_2}{\eta} + \mathbf{1}_{\{\delta_C(\rho) < 1\}} \frac{\delta_C(\rho)^2}{2} + \mathbf{1}_{\{\delta_C(\rho) \geq 1\}} \left(\delta_C(\rho) - \frac{1}{2} \right).$$

(i) We first prove that we may choose K_1 so that \underline{w} is a viscosity subsolution of the equation (5.5) satisfying the growth condition (5.6). Since \underline{w} has linear growth, δ_C is Lipschitz and vanishes at 0, we may choose $K_1 > 0$ such that $\underline{w}(\rho) \leq |\sigma\rho|^2/2$ for all $\rho \in \mathbb{R}^d$.

Moreover, \underline{w} is convex and has a gradient in the weak sense, which takes values in C , by definition of the support function. Then, for all $\rho_0 \in \mathbb{R}^d$, and $(p, X) \in J^{2,+}\underline{w}(\rho_0)$, we have $X \geq 0$, $p \in C$, and therefore:

$$\begin{aligned} & \max_{(i,j) \in \mathcal{J}} \left\{ -\frac{|\sigma^T \rho_0|^2}{2} - \frac{1}{2} \text{Tr} [\bar{\alpha} \bar{\alpha}^T X] + \eta \underline{w}(\rho_0), -\lambda^{i,j} + p_i - p_j \right\} \\ & \leq \max_{(i,j) \in \mathcal{J}} \left\{ -\frac{|\sigma^T \rho_0|^2}{2} + \eta \underline{w}(\rho_0), -\lambda^{i,j} + p_i - p_j \right\} \leq 0. \end{aligned}$$

Hence, \underline{w} is a viscosity subsolution.

(ii) We next prove that \bar{w} is a viscosity supersolution of the equation (5.5) satisfying the growth condition (5.6), for a convenient choice of K_2 . Consider arbitrary $\rho_0 \in \mathbb{R}^d$ and $(p, X) \in J^{2,-}\bar{w}(\rho_0)$.

Case 1: $\delta_C(\rho_0) < 1$, then ρ_0 must be bounded. Moreover, by definition, \bar{w} is Lipschitz. Hence, $p \in C$ and $X \geq 0$. In particular, p is bounded and by the definition of $J^{2,-}\bar{w}(\rho_0)$ so is X . We then have

$$-\frac{|\sigma^T \rho_0|^2}{2} - \frac{1}{2} \text{Tr} [\bar{\alpha} \bar{\alpha}^T X] + \eta \bar{w}(\rho_0) \geq -\frac{|\sigma^T \rho_0|^2}{2} - \frac{1}{2} \text{Tr} [\bar{\alpha} \bar{\alpha}^T X] + K_2.$$

Since ρ_0 and X are bounded, we may also choose K_2 large enough so that the above is positive. This implies the supersolution property in that case.

Case 2: $\delta_C(\rho_0) \geq 1$, then \bar{w} has a weak gradient which, by definition of the support function, takes values in ∂C , i.e. at least one of the gradient constraints in (5.5) is binding. This implies that the supersolution property is satisfied. \square

We next establish that \bar{w}^η is convex, by following the PDE argument of [21].

Lemma 5.1. *\bar{w}^η is convex and therefore is twice differentiable Lebesgue almost everywhere.*

Proof. 1. Let $\epsilon \in (0, 1)$, $\rho^0, \rho^1 \in \mathbb{R}^d$, $\bar{\rho} := (\rho^0 + \rho^1)/2$, and let us first prove that

$$\ell^\epsilon(\rho^0, \rho^1) := (1 - \epsilon) \bar{w}^\eta(\bar{\rho}) - (\bar{w}^\eta(\rho^0) + \bar{w}^\eta(\rho^1))/2 \longrightarrow -\infty \quad \text{as } |(\rho^0, \rho^1)| \longrightarrow \infty.$$

Let $(\rho_n^0, \rho_n^1) \in \mathbb{R}^d \times \mathbb{R}^d$ be such that $|\rho_n^0| + |\rho_n^1| \longrightarrow \infty$. Denote $\bar{\rho}_n := (\rho_n^0 + \rho_n^1)/2$. For large n , we have $\delta_C(\rho_n^0) + \delta_C(\rho_n^1) > 0$, and using the convexity of δ_C we also have

$$\begin{aligned} \Delta_n & := \frac{\ell^\epsilon(\rho_n^0, \rho_n^1)}{\delta_C(\rho_n^0) + \delta_C(\rho_n^1)} = (1 - \epsilon) \frac{\bar{w}^\eta(\bar{\rho}_n)}{\delta_C(\rho_n^0) + \delta_C(\rho_n^1)} - \frac{1}{2} \sum_{i=0,1} \frac{\delta_C(\rho_n^i)}{\delta_C(\rho_n^i) + \delta_C(\rho_n^{1-i})} \frac{\bar{w}^\eta(\rho_n^i)}{\delta_C(\rho_n^i)} \\ & \leq \frac{1 - \epsilon}{2} \frac{\bar{w}^\eta(\bar{\rho}_n)}{\delta_C(\bar{\rho}_n)} - \frac{1}{2} \sum_{i=0,1} \frac{\delta_C(\rho_n^i)}{\delta_C(\rho_n^i) + \delta_C(\rho_n^{1-i})} \frac{\bar{w}^\eta(\rho_n^i)}{\delta_C(\rho_n^i)}. \end{aligned}$$

Consider first the case where $(\delta_C(\bar{\rho}_n))_n$ is bounded, which is equivalent to the boundedness of $(|\bar{\rho}_n|)_n$ by (5.1). Then it is clear by the growth property (5.6) that $\Delta_n \rightarrow -\frac{1}{2} < 0$. Similarly, if $\delta_C(\bar{\rho}_n) \rightarrow \infty$, we see that $\limsup_{n \rightarrow \infty} \Delta_n \leq [(1 - \epsilon) - 1]/2 < 0$. In both case this proves the required result of this step.

2. For $\theta = (\rho^0, \rho^1, y^0, y^1) \in \mathbb{R}^{4d}$, set $\bar{\rho} := (\rho^0 + \rho^1)/2$, $\bar{y} := (y^0 + y^1)/2$, and define:

$$\varpi^\epsilon(\theta) := (1 - \epsilon)\bar{w}^n(\bar{\rho}) - (\bar{w}^n(y^0) + \bar{w}^n(y^1))/2, \quad \phi_n(\theta) := n \left(|\rho^0 - y^0|^2 + |\rho^1 - y^1|^2 \right) / 2.$$

Since \bar{w}^n is L -Lipschitz, we have

$$(\varpi^\epsilon - \phi_n)(\theta) = (1 - \epsilon)[(\bar{w}^n(\bar{\rho}) - \bar{w}^n(\bar{y}))] - \phi_n(\theta) + \ell^\epsilon(y^0, y^1) \leq \ell^\epsilon(y^0, y^1) + L^2/n.$$

By the first step, this shows that $(\varpi^\epsilon - \phi_n)(\theta) \rightarrow -\infty$ as $|\theta| \rightarrow \infty$, and that there is a global maximizer $\theta_n := (\rho_n^0, \rho_n^1, y_n^0, y_n^1)$ of the difference $\varpi^\epsilon - \phi_n$. Using then Crandall-Ishii's lemma and arguing exactly as in the proof of Proposition 3.1 (see also the proof of Lemma 3.7 in [21]), we obtain that for all (ρ^0, ρ^1)

$$\begin{aligned} (1 - \epsilon)\bar{w}^n(\bar{\rho}) - (\bar{w}^n(\rho^0) + \bar{w}^n(\rho^1))/2 &\leq (1 - \epsilon)\bar{w}^n(\bar{\rho}_n) - (\bar{w}^n(y_n^0) + \bar{w}^n(y_n^1))/2 \\ &\leq (2|\sigma^T \bar{\rho}_n|^2 - |\sigma^T y_n^0|^2 - |\sigma^T y_n^1|^2)/(4\eta). \end{aligned} \quad (5.8)$$

Following the same arguments as in the proof of Proposition 3.1, we next show that the sequence $(\theta_n)_n$ is bounded, and that there is a subsequence which converges to some $\theta_\epsilon := (\rho_\epsilon^0, \rho_\epsilon^1, y_\epsilon^0, y_\epsilon^1)$. The averages $\bar{\rho}_\epsilon$ and \bar{y}_ϵ are introduced similarly. Passing to the limit along this subsequence in (5.8), we obtain for all (ρ^0, ρ^1) :

$$\eta \left((1 - \epsilon)\bar{w}^n(\bar{\rho}) - [\bar{w}^n(\rho^0) + \bar{w}^n(\rho^1)]/2 \right) \leq \left(2|\sigma^T \bar{\rho}_\epsilon|^2 - |\sigma^T \rho_\epsilon^0|^2 - |\sigma^T \rho_\epsilon^1|^2 \right) \leq 0.$$

The proof is completed by sending ϵ to 0. □

As a consequence of the convexity of \bar{w}^n , we have the following result.

Lemma 5.2. *There is a constant $M > 0$ independent of η such that $J^{1,-}\bar{w}^n(\rho) \subset \partial C$, for all $\rho \in B_0(M)^c$.*

Proof. Let K_2 be the constant in Proposition 5.1. Since δ_C has linear growth, we can choose M large enough so that

$$K_2 + \delta_C(y) < \frac{|\sigma^T y|^2}{2} \quad \text{for all } |y| \geq M.$$

Fix $\rho \in \mathbb{R}^d$ satisfying $|\rho| \geq M$. Then, by the convexity of \bar{w}^n , $p \in J^{1,-}\bar{w}^n(\rho)$ if and only if p belongs to the subdifferential of \bar{w}^n at ρ . This implies, in particular, that $p \in C$. Furthermore, we have $(p, 0) \in J^{2,-}\bar{w}^n(\rho)$. Then, the supersolution property of \bar{w}^n yields

$$\max_{(i,j) \in \mathcal{J}} \left\{ -\frac{|\sigma^T \rho|^2}{2} + \eta \bar{w}^n(\rho), -\lambda^{i,j} + p_i - p_j \right\} \geq 0.$$

By the definition of M , we have $-\frac{|\sigma^T \rho|^2}{2} + \eta \bar{w}^n(\rho) \leq -\frac{|\sigma^T \rho|^2}{2} + K + \delta_C(\rho) < 0$, and therefore $\max_{(i,j) \in \mathcal{J}} \{-\lambda^{i,j} + p_i - p_j\} \geq 0$. Since $p \in C$, this means that this quantity is actually equal to zero, implying that $p \in \partial C$. \square

The following result is similar to Lemma 3.10 in [21].

Lemma 5.3. *For Lebesgue almost every $\rho \in \mathbb{R}^d$, we have $0 \leq D^2 \bar{w}^n(\rho) \leq \frac{1}{\eta} \|\sigma \sigma^T\|$.*

Proof. We fix $\epsilon \in (0, 1)$ and $z \in \mathbb{R}^d$ such that $0 < |z| < M$ which we will send to zero later. Set

$$\psi(\rho) := (1 - \epsilon)(\bar{w}^n(\rho + z) + \bar{w}^n(\rho - z)) - 2\bar{w}^n(\rho).$$

Since \bar{w}^n is Lipschitz,

$$\psi(\rho) \leq 2(1 - \epsilon)L|z| - 2\bar{w}^n(\rho) \leq 2(1 - \epsilon)LM - 2\bar{w}^n(\rho).$$

In view of the growth condition (5.6), as ρ approaches to infinity $\psi(\rho)$ tends to minus infinity. Let

$$\beta(\rho_1, \rho_2, \rho_3) := (1 - \epsilon)(\bar{w}^n(\rho_1 + z) + \bar{w}^n(\rho_2 - z)) - 2\bar{w}^n(\rho_3),$$

and

$$\phi_v(\rho_1, \rho_2, \rho_3) := \frac{1}{2v} \left(|\rho_1 - \rho_3|^2 + |\rho_2 - \rho_3|^2 \right).$$

We then have, again by the Lipschitz property of \bar{w}^n , that

$$\begin{aligned} (\beta - \phi_v)(\rho_1, \rho_2, \rho_3) &= (1 - \epsilon) (\bar{w}^n(\rho_1 + z) - \bar{w}^n(\rho_3 + z) + \bar{w}^n(\rho_2 + z) - \bar{w}^n(\rho_3 - z)) \\ &\quad + \psi(\rho_3) - \frac{1}{2v} \left(|\rho_1 - \rho_3|^2 + |\rho_2 - \rho_3|^2 \right) \\ &\leq L^2 v + \psi(\rho_3) \rightarrow -\infty, \quad \text{as } |\rho| \rightarrow \infty. \end{aligned}$$

Hence there exists $(\rho_1^v, \rho_2^v, \rho_3^v)$ which maximizes $\beta - \phi_v$. By applying Crandall-Ishii's lemma, for every $\rho > 0$, we can find symmetric matrices $(X, Y) \in \mathbb{S}^{2d} \times \mathbb{S}^d$ such that

$$\begin{aligned} (D_{\rho_1} \phi_v(\rho_1^v, \rho_2^v, \rho_3^v), D_{\rho_2} \phi_v(\rho_1^v, \rho_2^v, \rho_3^v), X) &\in \bar{J}^{2,+} (1 - \epsilon) (\bar{w}^n(\rho_1^v + z) + \bar{w}^n(\rho_2^v - z)) \\ (-D_{\rho_3} \phi_v(\rho_1^v, \rho_2^v, \rho_3^v), Y) &\in \bar{J}^{2,-} 2\bar{w}^n(\rho_3^v), \end{aligned}$$

and

$$\begin{pmatrix} X & 0 & 0 \\ 0 & -Y & 0 \\ 0 & 0 & 0 \end{pmatrix} \leq A + \rho A^2, \quad (5.9)$$

where

$$A := \begin{pmatrix} D^2\phi_v(\rho_1^v, \rho_2^v, \rho_3^v) & 0 \\ 0 & 0 \end{pmatrix} = \frac{1}{v} \begin{pmatrix} I_d & 0 & -I_d & 0 \\ 0 & I_d & -I_d & 0 \\ -I_d & -I_d & 2I_d & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

For $(X_1, X_2, X_3) \in \mathbb{S}^d \times \mathbb{R}^{d \times d} \times \mathbb{S}^d$, set $X := \begin{pmatrix} X_1 & X_2 \\ X_2^T & X_3 \end{pmatrix}$. Then, (5.9) implies that

$$X \leq \begin{pmatrix} -Y & 0 \\ 0 & 0 \end{pmatrix},$$

and in particular, $\text{Tr}[X_1 + X_3 - Y] \leq 0$.

We directly calculate that

$$\begin{aligned} \left(\frac{\rho_1^v - \rho_3^v}{v}, X_1 \right) &\in \bar{J}^{2,+}(\rho_1 \mapsto (1 - \epsilon)(\bar{w}^\eta(\rho_1 + z) + \bar{w}^\eta(\rho_2^v - z)))_{\rho_1 = \rho_1^v} \\ \left(\frac{\rho_2^v - \rho_3^v}{v}, X_3 \right) &\in \bar{J}^{2,+}(\rho_2 \mapsto (1 - \epsilon)(\bar{w}^\eta(\rho_1^v + z) + \bar{w}^\eta(\rho_2 - z)))_{\rho_2 = \rho_2^v} \\ \left(\frac{\rho_1^v + \rho_2^v - 2\rho_3^v}{v}, Y \right) &\in \bar{J}^{2,-}2\bar{w}^\eta(\rho_3^v). \end{aligned}$$

Since \bar{w}^η is a viscosity subsolution of (2.7), we deduce that for all $0 \leq i, j \leq d$

$$-\lambda^{j,i} \leq \frac{\rho_{1,i}^v - \rho_{3,i}^v}{(1 - \epsilon)v} - \frac{\rho_{1,j}^v - \rho_{3,j}^v}{(1 - \epsilon)v} \leq \lambda^{i,j},$$

and

$$-\lambda^{j,i} \leq \frac{\rho_{2,i}^v - \rho_{3,i}^v}{(1 - \epsilon)v} - \frac{\rho_{2,j}^v - \rho_{3,j}^v}{(1 - \epsilon)v} \leq \lambda^{i,j}.$$

From this we deduce that for all $0 \leq i, j \leq d$

$$-\lambda^{j,i} \leq \frac{\rho_{1,i}^v + \rho_{2,i}^v - 2\rho_{3,i}^v}{2\epsilon v} - \frac{\rho_{1,j}^v + \rho_{2,j}^v - 2\rho_{3,j}^v}{2(1 - \epsilon)v} \leq \lambda^{i,j}.$$

Since $\epsilon \in (0, 1)$, this implies that $\frac{\rho_1^v + \rho_2^v - 2\rho_3^v}{2v} \in \text{int}(C)$. Also $\frac{\rho_1^v + \rho_2^v - 2\rho_3^v}{2v} \in \bar{J}^{2,-}\bar{w}^\eta(\rho_3^v)$. Given that \bar{w}^η is a viscosity supersolution, we deduce that

$$-\frac{|\sigma^T \rho_3^v|^2}{2} - \frac{1}{2} \text{Tr} \left[\bar{\alpha} \bar{\alpha}^T \frac{Y}{2} \right] + \eta \bar{w}^\eta(\rho_3^v) \geq 0.$$

Since \bar{w}^η is a viscosity subsolution,

$$-\frac{|\sigma^T(\rho_1^v + z)|^2}{2} - \frac{1}{2(1 - \epsilon)} \text{Tr} [\bar{\alpha} \bar{\alpha}^T X_1] + \eta \bar{w}^\eta(\rho_1^v + z) \leq 0,$$

and

$$-\frac{|\sigma^T(\rho_2^v - z)|^2}{2} - \frac{1}{2(1-\epsilon)} \text{Tr}[\bar{\alpha}\bar{\alpha}^T X_3] + \eta\bar{w}^\eta(\rho_2^v - z) \leq 0.$$

Summing up the last three inequalities, we obtain that for all $\rho \in \mathbb{R}^d$,

$$\begin{aligned} (1-\epsilon)(\bar{w}^\eta(\rho+z) + \bar{w}^\eta(\rho-z)) - 2\bar{w}^\eta(\rho) &\leq (1-\epsilon)(\bar{w}^\eta(\rho_1^v+z) \\ &\quad + \bar{w}^\eta(\rho_2^v-z)) - 2\bar{w}^\eta(\rho_3^v) \\ &\leq \frac{1}{2\eta} \text{Tr}[\bar{\alpha}\bar{\alpha}^T(X_1+X_3-Y)] + \frac{1-\epsilon}{2\eta} \left(|\sigma^T(\rho_1^v+z)|^2 + |\sigma^T(\rho_2^v-z)|^2 \right) - \frac{|\sigma^T\rho_3^v|^2}{\eta} \\ &\leq \frac{1}{2\eta} \left(|\sigma^T(\rho_1^v+z)|^2 + |\sigma^T(\rho_2^v-z)|^2 \right) - \frac{|\sigma^T\rho_3^v|^2}{\eta}. \end{aligned} \quad (5.10)$$

We argue as in the proofs of Proposition 3.1 and Lemma 5.1, to show that $(\rho_1^v, \rho_2^v, \rho_3^v)$ is bounded and that there is a subsequence and a vector ρ^* such that they all converge to ρ^* along this subsequence. Using this result in (5.10), we obtain

$$\begin{aligned} \eta((1-\epsilon)(\bar{w}^\eta(\rho+z) + \bar{w}^\eta(\rho-z)) - 2\bar{w}^\eta(\rho)) &\leq \frac{|\sigma^T(\rho^*+z)|^2 + |\sigma^T(\rho^*-z)|^2}{2} - |\sigma^T\rho^*|^2 \\ &\leq \|\sigma\sigma^T\| |z|^2, \end{aligned}$$

where we used the mean value theorem and the fact that the Hessian matrix of the map $\rho \mapsto |\sigma^T\rho|^2/2$ is $\sigma\sigma^T$.

We finally let ϵ go to 0^+ , divide the inequality by $|z|^2$ and let $|z|$ go to zero to obtain the result, since we know that the second derivative of \bar{w}^η exists almost everywhere. \square

Corollary 5.2.

- (i) $\bar{w}^\eta \in C^{1,1}(\mathbb{R}^d)$.
- (ii) The set

$$\mathcal{C}_\eta := \{\rho \in \mathbb{R}^d, D\bar{w}^\eta(\rho) \in \text{int}(C)\}.$$

is open and bounded independently of $0 < \eta \leq 1$.

- (iii) $\bar{w}^\eta \in C^\infty(\mathcal{C}_\eta)$.
- (iv) There exists a constant $L > 0$ independent of η such that

$$0 \leq D^2\bar{w}^\eta(\rho) \leq L, \quad \rho \in \mathcal{C}_\eta.$$

Remark 5.2. Now that we have obtained some regularity for the function \bar{w}^η , we could hope to prove that it is indeed the value function of an infinite horizon stochastic control problem by means of a verification argument. However, the problem here is that the verification argument needs to identify an optimal control, which we cannot do in the present setting. Indeed, we do not know whether the solution to the reflexion problem on the free boundary defined by PDE (2.7) has a

solution, since we know nothing about the regularity of the free boundary. Notice that such a regularity was assumed in [28] in order to construct an optimal control.

Proof. The first result is a simple consequence of the previous Lemma 5.3. For the second one, the fact that \mathcal{O}_η is bounded independently of η follows from Lemma 5.2, and it is open because it is the inverse image of the open interval $(-\infty, 0)$ by the continuous application

$$\rho \mapsto \max_{(i,j) \in \mathcal{J}} \left\{ -\lambda^{i,j} + \frac{\partial \bar{w}^\eta}{\partial \rho_i}(\rho) - \frac{\partial \bar{w}^\eta}{\partial \rho_j}(\rho) \right\}.$$

Then the third result follows from classical regularity results for linear elliptic PDEs (see Theorem 6.17 in [18]), since we have on \mathcal{O}^η

$$-\frac{|\sigma^T \rho|^2}{2} - \frac{1}{2} \text{Tr} [\tilde{\alpha} \tilde{\alpha}^T D^2 \bar{w}^\eta(\rho)] + \eta \bar{w}^\eta(\rho) = 0.$$

Finally, by convexity of \bar{w}^η , we have for any ξ such that $|\xi| = 1$ and for any $\rho \in \mathcal{O}_\eta$

$$\xi^T D^2 \bar{w}^\eta(\rho) \xi \leq C_0 \text{Tr} [\tilde{\alpha} \tilde{\alpha}^T D^2 \bar{w}^\eta(\rho)] = 2C_0 \left(\eta \bar{w}^\eta(\rho) - \frac{|\sigma^T \rho|^2}{2} \right) \leq 2C_0(K_2 + \delta_C(\rho)),$$

and the result is a consequence of the fact that \mathcal{O}_η is bounded. \square

In the following result, we extend Proposition 3.12 of [21] to our context and show that \bar{w}^η is characterized by its values in $\overline{\mathcal{O}_\eta}$. The proof is very similar to the one given in [21]. We provide it here for the convenience of the reader.

Proposition 5.2. *We have for all $\rho \in \mathbb{R}^d$, $\bar{w}^\eta(\rho) = \inf_{y \in \overline{\mathcal{O}_\eta}} \{\bar{w}^\eta(y) + \delta_C(\rho - y)\}$.*

Proof. First of all, since $\bar{w}^\eta \in C^{1,1}(\mathbb{R}^d)$, it is easy to see by the mean value Theorem and the fact that $D\bar{w}^\eta \in C$ that for all $(\rho, y) \in \mathbb{R}^d \times \mathbb{R}^d$

$$-\delta_C(y - \rho) \leq \bar{w}^\eta(\rho) - \bar{w}^\eta(y) \leq \delta_C(\rho - y).$$

Hence we have $\bar{w}^\eta(y) + \delta_C(\rho - y) \geq \bar{w}^\eta(\rho)$. This implies that for $\rho \in \overline{\mathcal{O}_\eta}$ we have

$$\inf_{y \in \overline{\mathcal{O}_\eta}} \{\bar{w}^\eta(y) + \delta_C(\rho - y)\} = \bar{w}^\eta(\rho).$$

It remains to show that the result also holds for $\rho \in \overline{\mathcal{O}_\eta}^c$. Notice first that by convexity of \bar{w}^η , the minimum in the formula is necessarily achieved on $\partial \mathcal{O}_\eta$. Define then

$$\tilde{w}^\eta(\rho) := \inf_{y \in \partial \mathcal{O}_\eta} \{\bar{w}^\eta(y) + \delta_C(\rho - y)\}.$$

Let us then consider the following PDE

$$\max_{(i,j) \in \mathcal{F}} \left\{ -\lambda^{i,j} + \frac{\partial v}{\partial \rho_i}(\rho) - \frac{\partial v}{\partial \rho_j}(\rho) \right\} = 0, \quad \rho \in \overline{\mathcal{O}}_\eta^c$$

$$v(\rho) = \overline{w}^\eta(\rho), \quad \rho \in \partial \mathcal{O}_\eta.$$

It is easy to adapt the proof of Proposition 3.1 to obtain that this PDE satisfies a comparison principle. Since \overline{w}^η clearly solves it, it is the unique solution. Then by the usual properties of inf-convolutions, it is clear that the function \tilde{w}^η which is convex, has a gradient in the weak sense which is in C . This implies that \tilde{w}^η is a subsolution of the above PDE. Moreover, since δ_C also has a gradient in the weak sense which is in ∂C (by convexity), using the fact that by properties of the inf-convolution of convex functions, the subgradient of \tilde{w}^η at any point is contained in a subgradient of δ_C , we have that \tilde{w}^η dominates every subsolution of the PDE which coincides with \overline{w}^η on $\partial \mathcal{O}_\eta$. By usual results of the theory of viscosity solutions (see [10]) this proves that \tilde{w}^η is also a supersolution. By uniqueness, we obtain the desired result. \square

Corollary 5.3. *For Lebesgue almost every $\rho \in \overline{\mathcal{O}}_\eta^c$, $D^2 \overline{w}^\eta(\rho) = 0$. Hence, the second derivative of \overline{w}^η is bounded almost everywhere, independently of η in the whole space \mathbb{R}^d .*

Proof. Let $\rho \in \overline{\mathcal{O}}_\eta^c$. By Proposition 5.2, there exists $y \in \partial \mathcal{O}_\eta$ such that $\overline{w}^\eta(\rho) = \overline{w}^\eta(y) + \delta_C(\rho - y)$. We then have for any $z \in \mathbb{R}^d$

$$\frac{\overline{w}^\eta(\rho + z) + \overline{w}^\eta(\rho - z) - 2\overline{w}^\eta(\rho)}{|z|^2} = \frac{\overline{w}^\eta(\rho + z) + \overline{w}^\eta(\rho - z) - 2\overline{w}^\eta(y) - 2\delta_C(\rho - y)}{|z|^2}$$

$$\leq \frac{\delta_C(\rho + z - y) + \delta_C(\rho - z - y) - 2\delta_C(\rho - y)}{|z|^2}.$$

Now, the function $\rho \rightarrow \delta_C(\rho)$ is continuous and clearly C^∞ almost everywhere (actually except on $d(d+1)$ hyperplanes which therefore have Lebesgue measure 0). More than that, this function is piecewise linear, which implies that its Hessian is null almost everywhere. By letting $|z|$ go to 0 above, we obtain the desired result. The last result is now a simple consequence of Corollary 5.2(iv). \square

Then, the uniform estimates obtained above allow us to prove our main existence result in Theorem 3.2.

Proof of Theorem 3.2. By the uniform estimates of Corollary 5.3, we may follow the arguments of Section 4.2 in [21] to construct a strictly positive sequence $(\eta_n)_{n \geq 0}$ converging to zero, $\bar{a} \in \mathbb{R}$ and $\bar{w} \in C^{1,1}(\mathbb{R}^d)$ such that

$$\lim_{n \rightarrow +\infty} \eta_n \overline{w}^{\eta_n}(\rho^{\eta_n}) = \bar{a}, \quad \text{and} \quad \overline{w}^{\eta_n} \xrightarrow[n \rightarrow +\infty]{} \bar{w} \text{ in } C_{loc}^1(\mathbb{R}^d),$$

where ρ^{η_n} is a global minimizer of \overline{w}^{η_n} . Corollary 5.2 implies that the limiting function \bar{w} satisfies the required properties. \square

A. Appendix

Proof of Lemma 4.2. We assume throughout the proof that the parameter ϵ is less than one. In view of the gradient constraints of the dynamic programming equation satisfied by v^ϵ , we know that for all $1 \leq i \leq d$, we have in the viscosity sense

$$\Lambda_{i,0}^\epsilon(v_x^\epsilon, v_y^\epsilon) \geq 0, \quad \Lambda_{0,i}^\epsilon(v_x^\epsilon, v_y^\epsilon) \geq 0. \quad (\text{A.1})$$

Now define

$$\hat{v}^\epsilon(s, z, \zeta) := v^\epsilon(s, z - \epsilon\zeta \cdot \mathbf{1}_d - \mathbf{y}(s, z) \cdot \mathbf{1}_d, \epsilon\zeta + \mathbf{y}(s, z)).$$

We directly calculate that (A.1) implies for all $1 \leq i \leq d$ in the viscosity sense

$$\epsilon^4 \lambda^{i,0} \hat{v}_z^\epsilon(s, z, \zeta) - \epsilon^3 \lambda^{i,0} \mathbf{y}_z^i(s, z) \cdot \hat{v}_\zeta^\epsilon(s, z, \zeta) + (1 + \epsilon^3 \lambda^{i,0}) \hat{v}_{\zeta_i}^\epsilon(s, z, \zeta) \geq 0, \quad (\text{A.2})$$

and

$$\epsilon^4 \lambda^{0,i} \hat{v}_z^\epsilon(s, z, \zeta) - \epsilon^3 \lambda^{0,i} \mathbf{y}_z^i(s, z) \cdot \hat{v}_\zeta^\epsilon(s, z, \zeta) - \hat{v}_{\zeta_i}^\epsilon \geq 0. \quad (\text{A.3})$$

Using the fact that all the $\mathbf{y}_z^i(s, z)$ are strictly positive (see Assumption 3.1), we can multiply (A.2) by $\mathbf{y}_z^i(s, z)$ and sum to obtain, once more in the viscosity sense

$$\left(1 - \epsilon^3 \sum_{i=1}^d \frac{\mathbf{y}_z^i(s, z) \lambda^{i,0}}{1 + \epsilon^3 \lambda^{i,0}}\right) \mathbf{y}_z^i(s, z) \cdot \hat{v}_\zeta^\epsilon(s, z, \zeta) \geq -\epsilon^4 \sum_{i=1}^d \frac{\lambda^{i,0}}{1 + \epsilon^3 \lambda^{i,0}} \hat{v}_z^\epsilon(s, z, \zeta). \quad (\text{A.4})$$

Now, we have by Assumption 3.1

$$1 - \epsilon^3 \sum_{i=1}^d \frac{\mathbf{y}_z^i(s, z) \lambda^{i,0}}{1 + \epsilon^3 \lambda^{i,0}} = 1 - \sum_{i=1}^d \mathbf{y}_z^i(s, z) + \sum_{i=1}^d \frac{\mathbf{y}_z^i(s, z)}{1 + \epsilon^3 \lambda^{i,0}} \geq 0.$$

Using this inequality in (A.4) yields, in the viscosity sense

$$\mathbf{y}_z(s, z) \cdot \hat{v}_\zeta^\epsilon(s, z, \zeta) \geq -\frac{\sum_{i=1}^d \frac{\lambda^{i,0}}{1 + \epsilon^3 \lambda^{i,0}}}{1 - \epsilon^3 \sum_{i=1}^d \frac{\mathbf{y}_z^i(s, z) \lambda^{i,0}}{1 + \epsilon^3 \lambda^{i,0}}} \epsilon^4 \hat{v}_z^\epsilon(s, z, \zeta). \quad (\text{A.5})$$

Plugging this estimate in (A.2) and (A.3), we obtain in the viscosity sense

$$\hat{v}_{\zeta_i}^\epsilon(s, z, \zeta) \leq \lambda^{0,i} \epsilon^4 \left(1 + \frac{\sum_{i=1}^d \frac{\lambda^{i,0}}{1 + \epsilon^3 \lambda^{i,0}}}{1 - \epsilon^3 \sum_{i=1}^d \frac{\mathbf{y}_z^i(s, z) \lambda^{i,0}}{1 + \epsilon^3 \lambda^{i,0}}}\right) \hat{v}_z^\epsilon(s, z, \zeta) \leq \epsilon^4 \bar{\lambda} \left(1 + \frac{\bar{\lambda} d}{c_0}\right) \hat{v}_z^\epsilon(s, z, \zeta),$$

and

$$\hat{v}_{\zeta_i}^\epsilon(s, z, \zeta) \geq -2\epsilon^4 \underline{\lambda} \hat{v}_z^\epsilon(s, z, \zeta).$$

By the concavity of v^ϵ , its gradient exists almost everywhere. Moreover, since we assumed that \mathbf{y} is smooth, this implies that \hat{v}_z^ϵ also exists almost everywhere. Hence, we conclude from the above estimates that

$$|\hat{v}_\xi^\epsilon| \leq A\epsilon^4 \hat{v}_z^\epsilon, \quad \text{where } A := \sqrt{d} \max \left\{ \bar{\lambda} \left(1 + \frac{\bar{\lambda}^2 d}{c_0} \right), 2\lambda \right\}. \quad (\text{A.6})$$

Using these estimates, we can now finish the proof exactly as in the proof of Lemma 6.2 in [40]. \square

Lemma A.1. *Let Assumption 3.4 hold. For any $m > 0$, the function w^m satisfies:*

(i) \tilde{w}^m is C^2 , convex in ξ , and we have for all $0 \leq i, j \leq d$

$$\begin{aligned} \tilde{w}_{\xi_i}^m(s, z, \xi) &= \int_{\mathbb{R}^d} k^m(\zeta) w_{\xi_i}(s, z, \xi - \zeta) d\zeta, \\ \tilde{w}_{\xi_i \xi_j}^m(s, z, \xi) &= \int_{\mathbb{R}^d} k^m(\zeta) w_{\xi_i \xi_j}(s, z, \xi - \zeta) d\zeta. \end{aligned}$$

Moreover, $0 \leq \tilde{w}^m(s, z, \xi) \leq L v_z(s, z)(1+m)(1+|\xi|)$.

(ii) \tilde{w}^m is smooth in (s, z) , and satisfies the following estimates, uniformly in m ,

$$\begin{aligned} (|\tilde{w}^m| + |\tilde{w}_s^m| + |\tilde{w}_{ss}^m| + |\tilde{w}_z^m| + |\tilde{w}_{sz}^m| + |\tilde{w}_{zz}^m|)(s, z, \xi) &\leq C(s, z)(1+m)(1+|\xi|) \\ (|\tilde{w}_\xi^m| + |\tilde{w}_{s\xi}^m| + |\tilde{w}_{z\xi}^m|)(s, z, \xi) &\leq C(s, z) \\ |\tilde{w}_{\xi\xi}^m(s, z, \xi)| &\leq C(s, z) \mathbf{1}_{\xi \in \bar{B}(s, z)}, \end{aligned} \quad (\text{A.7})$$

where $C(s, z)$ is a continuous function depending on the Merton value function and its derivatives, and $B(s, z)$ is some ball with continuous radius and center in (s, z) .

(iii) For every $0 \leq i, j \leq d$ and every (s, z, ξ)

$$-\lambda^{i,j} v_z(s, z) + \tilde{w}_{\xi_i}^m(s, z, \xi) - \tilde{w}_{\xi_j}^m(s, z, \xi) \leq 0.$$

(iv) For every (s, z, ξ) , we have

$$\frac{1}{2} v_{zz}(s, z) \int_{\mathbb{R}^d} k^m(x) |\sigma(s)(\xi - \zeta)|^2 d\zeta - \frac{1}{2} \text{Tr} [\alpha \alpha^T(s, z) \tilde{w}_{\xi\xi}^m(s, z, \xi)] + a(s, z) \leq 0.$$

We next construct a useful function which plays a major role in our subsequent proof.

Lemma A.2. *For any $\delta \in (0, 1)$ and any $\nu > 0$, there exists $a^\delta := a^{\delta, \nu} > 1$ and a function $h^{\delta, \nu} : \mathbb{R}^d \rightarrow [0, 1]$ such that $h^{\delta, \nu}$ is C^∞ , $h^{\delta, \nu} = 1$ on $\mathcal{B}_1(0)$ and $h^{\delta, \nu} = 0$ on $B_{a^\delta}(0)^c$. Moreover, for any $1 \leq i, j \leq d$ and for any $\xi \in \mathbb{R}^d$*

$$|h_{\xi_i}^{\delta, \nu}(\xi)| \leq \frac{\nu \delta}{3}, \quad |\xi| |h_{\xi_i}^{\delta, \nu}| \leq \nu \delta, \quad \text{and} \quad |\xi| |h_{\xi\xi}^{\delta, \nu}| + |h_{\xi\xi}^{\delta, \nu}| \leq C^*,$$

for some constant C^* independent of δ .

B. Numerical Results (By Loïc Richier and Bertrand Rondepierre)

In this section, we report some examples of numerical results. We follow the numerical scheme suggested in Campillo [8] which combines the finite differences approximation and the policy iteration method in order to produce an approximation of the solution of the first corrector equation. We recall that a first order approximation of the no-transaction region, for fixed $s \in (0, \infty)^d$, is deduced from the free boundary set \mathcal{C}_0 of Proposition 3.2 by:

$$\hat{\mathbf{N}}\mathbf{T}_\epsilon(s) := \{(x, y) \in \mathbb{R}^{d+1} : y = \theta(s, z) + \epsilon\eta(s, z)\mathcal{C}_0\}. \quad (\text{B.1})$$

Also, as explained in Remark 5.2 above, the first corrector equation corresponds to a singular ergodic control problem whose corresponding optimal control processes can be viewed as a first order approximation of the optimal transfers in the original problem of optimal investment and consumption under transaction costs.

For simplicity, the present numerical results are obtained in the two-dimensional case $d = 2$ for power utility with Black-Scholes dynamics. Under this simplification, it is well-known that the value function of the Merton problem v is homogeneous in z , which in turn implies that the optimal control $\mathbf{y}(z)$ is actually linear in z . Hence, the coefficient $\bar{\alpha}$ reduces to a constant and the solution of the first corrector equation is independent of z .

B.1. Overview of the Numerical Scheme

As explained above, we use the formal link established between the first corrector equation and the ergodic control problem to perform our numerical scheme. The first step in the numerical approximation for the first corrector equation is to transform the original ergodic control problem to a control problem for a Markov process in continuous time and finite state space. To do so, we restrict the domain of the variable ρ to a bounded (and large enough) subset of \mathbb{R}^2 , denoted by \mathcal{D} , which is then discretized with a regular grid containing N^2 points. Then, the diffusion part of the first corrector equation is approximated using a well-chosen finite difference scheme (we refer the reader to [8] for more details). The discretized HJB equation thus takes the following form:

$$\min_{m \geq 0} \left(\sum_{\rho' \in \mathcal{D}} \mathcal{L}_h^m(\rho, \rho') \bar{w}(\rho') + f^m(\rho) \right) = \bar{a}, \quad \text{for all } \rho \in \mathcal{D},$$

where \mathcal{L}_h^m is the discretized version of the infinitesimal generator appearing in the first order corrector equation, m is a 3×3 matrix that corresponds to the controls and

$$f^m(\rho) := \frac{|\sigma^T \rho|^2}{2} + \text{Tr}[\lambda^T m], \quad \lambda = (\lambda^{i,j})_{0 \leq i, j \leq 2}, \quad \rho \in \mathcal{D}.$$

The policy iteration algorithm corresponds now to the following iterative procedure which starts from an arbitrary initial policy m^0 , and involves the two following steps:

- (i) Given a policy m^j , we compute for all $\rho \in D$ the solution (\bar{w}^j, \bar{a}^j) of the linear system

$$\sum_{\rho' \in \mathcal{D}} \mathcal{L}_h^{m^j}(\rho, \rho') \bar{w}^j(\rho') + f^m(\rho) = \bar{a}^j.$$

Notice that this is a linear system of N^2 equations for the N^2 unknowns $(\bar{w}^j(\rho), \rho \in \mathcal{D} \setminus \{0\})$ and \bar{a}^j . Recall that $\bar{w}^j(0) = 0$ is given.

- (ii) We update the optimal control by solving the N^2 minimization problems

$$m^{j+1}(\rho) \in \arg \min_{m \geq 0} \sum_{\rho' \in \mathcal{D}} \{ \mathcal{L}_h^m(\rho, \rho') \bar{w}^j(\rho') + f^m(\rho) \}.$$

Finally, the algorithm is stopped whenever the difference between \bar{a}^{j+1} and \bar{a}^j is smaller than some initially fixed error.

B.2. The Numerical Results

We present below different figures obtained with the above numerical scheme, representing the domain \mathcal{C}_0 and partitioning the state space in different regions depending on which controls are active or not. We remind once more the reader that \mathcal{C}_0 provides a first order approximation (B.1) of the true no-transaction region. For later reference, we provide our color correspondence, where we remind the reader that 0 corresponds to the cash account and 1 and 2 to the two risky assets, and i/j indicates that a positive transfer from asset i to j , or from j to i , occurs in the corresponding region.

Transactions	NT	1/2	0/1	0/2	0/1 and 0/2	1/2 and 0/1	1/2 and 0/2	1/2 and 0/1 and 0/2
Color		Blue	Red	Yellow	Orange	Purple	Green	Black

B.2.1. Cash-to-Asset Only. In all existing literature addressing either numerical procedures or asymptotic expansions for the multidimensional transaction costs problem, the transactions are only allowed between a given asset and the bank account, which in our setting translates into $\lambda^{i,j} < +\infty$, if and only if $i = 0$ or $j = 0$, see for instance Muthuraman and Kumar [30] or Bichuch and Shreve [6]. In this first section, we restrict ourselves to this case and show that our numerical procedure reproduces the earlier findings.

First, we consider the following symmetric transaction costs structure with the following values for σ

$$\lambda_0 = \begin{pmatrix} 0 & 0.001 & 0.001 \\ 0.001 & 0 & \infty \\ 0.001 & \infty & 0 \end{pmatrix}, \quad \sigma_0 = I_d, \quad \sigma_- = \begin{pmatrix} 1 & -0.25 \\ -0.25 & 1 \end{pmatrix}, \quad \sigma_+ = \begin{pmatrix} 1 & 0.25 \\ 0.25 & 1 \end{pmatrix}.$$

As expected and in line with the results of [30], the no-transaction region in the uncorrelated case is a rectangle (see Figure 1). Then, under a possible correlation between the assets, the shape of the region is modified to a parallelogram, the direction of the deformation depending on the sign of the correlation.




Figure 1. Uncorrelated (left), negative correlation (middle), positive correlation (right).




Figure 2. Higher correlations: negative (left), positive (middle), asymmetric costs (right).

In the next Figure 2, the first two graphs keep the same transaction costs structure but with higher correlations induced by the volatility matrices σ_{--} and σ_{++} , while in the third one we isolate the impact of letting the transaction costs from one asset to the cash be higher than for the other asset by using σ_0 , and the transaction costs structure λ'

$$\sigma_{--} = \begin{pmatrix} 1 & -0.25 \\ -0.1 & 1 \end{pmatrix}, \quad \sigma_{++} = \begin{pmatrix} 1 & 0.25 \\ 0.1 & 1 \end{pmatrix}, \quad \lambda' = \begin{pmatrix} 0 & 0.001 & 0.002 \\ 0.001 & 0 & \infty \\ 0.002 & \infty & 0 \end{pmatrix}.$$

By comparison to the first figures, we observe that modifying the correlation induces a rotation of the no-transaction regions, while in the asymmetric costs case, we still observe a rectangle but with modified dimensions (see Figure 3). More precisely, transactions between the first asset and the cash account occur more often since they are cheaper.

B.2.2. Possible Transfers Between all Assets. In this section we allow for transactions between all assets, a feature which was not considered in any of the existing numerical approximations in the literature on the present problem. We start by fixing a symmetric transaction costs structure λ_0 , and we illustrate the impact of correlation by considering the volatility matrices σ_0 , σ_- and σ_+ . We also report a figure obtained with the same parameters, but with more precise computations and without color.

Our first observation is that transactions between the two assets do occur, and more importantly that as a consequence, the no-transaction region seems to no longer be convex, an observation which, as far as we know, was not made before in the literature. Moreover, as in the previous section, the introduction of correlation induces a deformation of the no-transaction region.




Figure 3. Uncorrelated (upper left), negative correlation (upper right), positive correlation (lower left), precise result (lower right).




Figure 4. Asymmetric transaction costs.

Figure 4 shows the impact of an asymmetric transaction costs structure. The volatility matrix is set to σ_0 and the transaction costs matrix to λ' . As expected, there are almost no transactions between the cash account and asset 1, since they are

twice as expensive as the other ones. Surprisingly, we also observe the occurrence of small zones (in black and violet), where transactions are simultaneously performed between the assets and between the assets and the cash account.

Funding

The second author's research was partly supported by the European Research Council under the grant 228053-FiRM, the Swiss Finance Institute and by the ETH Foundation. The third author's research was supported by the Chair Financial Risks of the Risk Foundation sponsored by Société Générale, and the Chair Finance and Sustainable Development sponsored by EDF and Calyon.

References

- [1] Atkinson, C., Mokkavesa, S. (2004). Multi-asset portfolio optimization with transaction cost. *App. Math. Finance* 11:95–123.
- [2] Barles, G., Perthame, B. (1987). Discontinuous solutions of deterministic optimal stopping problems. *Math. Modeling Numerical Analysis* 21:557–579.
- [3] Barles, G., Soner, H.M (1998). Option pricing with transaction costs and a nonlinear Black-Scholes equation. *Finance and Stochastics* 2:369–397.
- [4] Benes, V.E., Shepp, L.A., Witsenhausen, H.S. (1980). Some solvable stochastic control problems. *Stochastics* 4:39–83.
- [5] Bichuch, M. (2011). Asymptotic analysis for optimal investment in finite time with transaction costs. Preprint.
- [6] Bichuch, M., Shreve, S.E. (2012). Utility maximization trading two futures with transaction costs. *SIAM J. Fin. Math.* 3:433–458.
- [7] Caffarelli, L.A., Souganidis, P.E. (2010). Rates of convergence for the homogenization of fully nonlinear uniformly elliptic pde in random media. *Inventiones Mathematicae* 180:301–360.
- [8] Campillo, F. (1995). Optimal ergodic control of nonlinear stochastic systems. In Krée, P., Wedig, W., eds., *Probabilistic Methods in Applied Physics*. Berlin: Springer-Verlag, pp. 239–269.
- [9] Constantinides, G.M. (1986). Capital market equilibrium with transaction costs. *J. Pol. Econ.* 94:842–862.
- [10] Crandall, M.G., Ishii, H., Lions, P.L. (1992). User's guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc.* 27:1–67.
- [11] Davis, M.H.A., Norman, A. (1990). Portfolio selection with transaction costs. *Math. O.R.* 15:676–713.
- [12] Davis, M.H.A, Panas, V.G., Zariphopoulou, T., (1993) European option pricing with transaction costs. *SIAM Journal on Control and Optimization* 31:470–493
- [13] Dumas, B., Luciano, E. (1991). An exact solution to a dynamic portfolio choice problem under transaction costs. *J. Finance* 46:577–595.
- [14] Evans, L.C. (1989). The perturbed test function technique for viscosity solutions of partial differential equations. *Proc. Royal Soc. Edinburgh Sect. A* 111:359–375.
- [15] Evans, L.C. (1992). Periodic homogenization of certain fully nonlinear partial differential equations. *Proc. Royal Soc. Edinburgh Sec. A Math.* 120:245–265.
- [16] Fleming, W.H., Soner, H.M. (1989). Asymptotic expansions for Markov processes with Levy generators. *Appl. Math. Optim.* 19:203–223.

- [17] Fleming, W.H., Souganidis, P.E. (1986). Asymptotic series and the method of vanishing viscosity. *Indiana Univ. Math. J.* 35:425–447.
- [18] Gilbarg, D., Trudinger, N. (1998). *Elliptic Partial Differential Equations of Second Order*. Berlin: Springer-Verlag.
- [19] Gerhold, S., Muhle-Karbe, J., Schachermayer, W. (2012). Asymptotics and duality for the Davis and Norman problem. *Stochastics* 84:625–641.
- [20] Goodman, J., Ostrov, D. (2010). Balancing Small Transaction Costs with Loss of Optimal Allocation in Dynamic Stock Trading Strategies. *SIAM J. Appl. Math.* 70:1977–1998.
- [21] Hynd, R. (2012). The eigenvalue problem of singular ergodic control. *Comm. Pure. Appl. Math.* 65:649–682.
- [22] Hynd, R. (2013). Analysis of Hamilton-Jacobi-Bellman equations arising in stochastic singular control. *ESAIM Cont. Optim. Calc. Var.* 19:112–128.
- [23] Janecek, K., Shreve, S.E. (2004). Asymptotic analysis for optimal investment and consumption with transaction costs. *Finance and Stochastics* 8:181–206.
- [24] Kabanov, Y., Safarian, M. (2009). *Markets with Transaction Costs*. Berlin: Springer-Verlag.
- [25] Lehoczky, J., Sethi, S.P., Soner, H.M., Taksar, M.I. (1991). An asymptotic analysis of hierarchical control of manufacturing systems under uncertainty. *Math. Oper. Res.* 16:596–608.
- [26] Lions, P.L., Souganidis, P.E. (2005). Homogenization of degenerate second-order PDE in periodic and almost periodic environments and applications. *Annales de L'Institut Henri Poincaré - Analyse Non Lineaire* 22:667–677.
- [27] Magill, M.J.P., Constantinides, G.M. (1976). Portfolio selection with transaction costs. *J. Econ. Theory* 13:245–263.
- [28] Menaldi, J.L., Robin, M., Taksar, M.I. (1992). Singular ergodic control for multidimensional Gaussian processes. *Math. Control Signals Systems* 5:93–114.
- [29] Merton, R.C. (1990). *Continuous Time Finance*. Oxford: Blackwell Publishing Ltd.
- [30] Muthuraman, K., Kumar, S. (2006). Multidimensional portfolio optimization with proportional transaction costs. *Math. Fin.* 16:301–335.
- [31] Papanicolaou, G., Varadhan, S.R.S. (1979). Boundary value problems with rapidly oscillating random coefficients. In: *Proceedings of Conference on Random Fields, Esztergom, Hungary, 1979; Seria Colloquia Mathematica Societatis Janos Bolyai*, Vol. 27. New York: North Holland, pp. 835–873.
- [32] Possamai, D., Soner, H.M., Touzi, N. (2012). Large liquidity expansion of super-hedging costs. *Aysmptotic Analysis*, 79:45–64.
- [33] Rogers, L.C.G., (2004). Why is the effect of proportional transaction costs $O(2/3)$? In: Yin, G., Zhang, Q., eds., *Mathematics of Finance*. AMS Contemporary Mathematics Series 351, Providence, RI: American Mathematical Society, pp. 303–308.
- [34] Sethi, S., Soner, H.M., Zhang, Q., Jiang, J. (1992). Turnpike Sets and Their Analysis in Stochastic Production Planning Problems. *Math. Op. Res.* 17: 932–950.
- [35] Shreve, S.E., Soner, H.M. (1989). Regularity of the value function of a two-dimensional singular stochastic control problem. *SIAM J. Cont. Opt.* 27:876–907.
- [36] Shreve, S.E., Soner, H.M. (1991). A free boundary problem related to singular stochastic control: parabolic case. *Comm. Part. Diff. Eqs.* 16:373–424.

- [37] Shreve, S.E., Soner, H.M. (1994). Optimal investment and consumption with transaction costs. *Ann. Appl. Prob.* 4:609–692.
- [38] Soner, H.M. (1993). Singular perturbations in manufacturing. *SIAM J. Control Opt.* 31:132–146.
- [39] Soner, H.M., Shreve, S.E., Cvitanic, J. (1995). There is no nontrivial hedging portfolio for option pricing with transaction costs. *Ann. Appl. Prob.* 5:327–355.
- [40] Soner, H.M., Touzi, N. (2013). Homogenization and asymptotics for small transaction costs. *SIAM J. Control Opt.* 51:2893–2921.
- [41] Souganidis, P.E., (1999) Stochastic homogenization for Hamilton-Jacobi equations and applications. *Asymptot. Anal.* 20:1–11.
- [42] Whalley, A.E., Willmott, P. (1997). An asymptotic analysis of the Davis, Panas & Zariphopoulou model for option pricing with transaction costs. *Math. Finance* 7:307–324.